

# **Syphilis Screening During Pregnancy: New York State Laws and Regulations**

---

## **Frequently Asked Questions for Providers**



**Department  
of Health**



## General

### 1. Q: When during pregnancy is syphilis screening required?

- A:** All pregnant persons are required to be screened<sup>1</sup> serologically for syphilis up to three times during pregnancy:
- At the time pregnancy is first diagnosed<sup>2</sup>
  - At 28 weeks of pregnancy, or as soon thereafter as is reasonably possible, but no later than at 32 weeks of pregnancy<sup>3</sup>
  - Again at delivery<sup>4</sup>

Practitioners are also encouraged to combine the third trimester syphilis screening with the recommended third trimester HIV screening per New York State Department of Health AIDS Institute Clinical Guidelines.

### 2. Q: What are the specific requirements for screening at delivery? Is this a screening for the birthing parent, the infant, and/or the cord blood?

- A:** The recommended practice is to screen the pregnant person for syphilis on admission to delivery using standard serologic testing. If the pregnant person screens positive for syphilis, the baby/infant shall be tested as well. Alternatively, if the provider is unable to screen the pregnant person on admission or immediately following delivery, per the most recent guidance from CDC it is best to test the infant body blood because umbilical cord blood can become contaminated with maternal blood and yield a false-positive result for non-treponemal (lipoidal antigen) tests, and Wharton's jelly within the umbilical cord can yield a false-negative result. The same nontreponemal (lipoidal antigen) test should be used for the infant that was used for the mother at delivery so titer levels can be compared.

---

<sup>1</sup> The Centers for Disease Control and Prevention recommends serologic testing for syphilis, including both nontreponemal and treponemal tests, particularly for high-risk populations and symptomatic individuals. <https://www.cdc.gov/mmwr/volumes/73/rr/pdfs/rr7301a1-H.pdf>

<sup>2</sup> New York State Public Health Law §2308 (the phrase "at the time of first examination" is interpreted to mean "at the time pregnancy is first diagnosed")

<sup>3</sup> New York State Public Health Law §2308, effective May 3, 2024

<sup>4</sup> New York State Public Health Law §§2500-a, 2308; 10 New York Codes, Rules and Regulations (NYCRR) §69-2.2

As indicated in the regulation associated with syphilis screening at delivery, the cord blood test requirement is waived if the infant body blood is tested.

### **3. Q: When did syphilis screening requirements go into effect?**

**A: Syphilis screening at first exam:** The requirement to submit the initial blood sample taken at the first exam has been a requirement in New York State since 1953.

**Syphilis screening during the third trimester:** Effective May 3, 2024, a syphilis screen is required during the third trimester of pregnancy. This is an additional screening requirement due to an increase in pregnant persons acquiring syphilis later in pregnancy.

**Syphilis screening at delivery:** In December 1989, a requirement to screen the cord blood of infants for syphilis was added to the New York State health regulations (10 NYCRR). In practice, this requirement has been implemented as a screening for syphilis at delivery. The reasoning here is two-fold: 1) the law states that the cord blood screening requirement is waived if body blood from the pregnant person is screened for syphilis at the time of birth, so long as the infant's body blood is tested after any positive test result of the pregnant person's blood, and 2) cord blood testing has been associated with both false positive and false negative results.<sup>5</sup>

### **4. Q: Did the third trimester screening law change from screening at the time of delivery to third trimester screening?**

**A:** No. As per Public Health Law Section §2308, in addition to screening at the time of first examination, every practitioner attending pregnant persons shall screen for syphilis during the third trimester of pregnancy consistent with guidance and regulations of the New York State Department of Health. There has been no change to the requirement in 10 NYCRR §69-2.2 to perform a syphilis screen again at delivery. Please see New York State's [interim guidance](#) for more information.

---

<sup>5</sup> Papp JR, Park IU, Fakile Y, Pereira L, Pillay A, Bolan GA. CDC Laboratory Recommendations for Syphilis Testing, United States, 2024. MMWR Recomm Rep 2024;73(No. RR-1):1–32. DOI: <https://stacks.cdc.gov/view/cdc/147800>

**5. Q. Who is required to do the syphilis screening during pregnancy?**

**A:** Every physician or other authorized practitioner attending pregnant persons in New York State.

**6. Q: What if the pregnant person in my care states they have only one sexual partner, or they have not had sexual intercourse since their first trimester screen, and are therefore not at risk of syphilis?**

**A:** Syphilis screening requirements are not waived for pregnant persons who report one sexual partner, or for persons who have reported not being sexually active since the first trimester screening. However, it should be noted that for persons reporting only one sexual partner, transmission is still possible.

A patient centered approach to this sexual health conversation is recommended to promote a productive discussion and share decision making with the patient.

**7. Q: What if the pregnant person refuses syphilis screening during their pregnancy?**

**A:** The law requires practitioners to submit the initial blood sample taken at the first exam for serological screening for syphilis and a syphilis screen during the third trimester. The law is intended to detect late seroconversion in pregnancy. If the patient refuses syphilis screening, document the refusal in the medical record.

**8. Q: Are there other syphilis screening requirements if a person delivers a stillborn infant?**

**A:** The New York regulation at 10 NYCRR §69-2.2 applies when an infant is born alive or dead after 22 weeks gestation; however, any person who delivers a stillborn infant after 20 weeks gestation should be screened for syphilis.<sup>6</sup>

---

<sup>6</sup> <https://www.cdc.gov/std/treatment-guidelines/syphilis-pregnancy.htm>

**9. Q: Why are these syphilis screening requirements in place?**

**A: Syphilis screening at first exam/visit:** This requirement is in place to promote a healthy start to a pregnancy, and to ensure the pregnant person is adequately treated prior to delivery. Early identification and treatment of syphilis is the best way to prevent transmission to the unborn infant, and to ensure the health status of the pregnant person.

**Syphilis screening during the third trimester:** For individuals who screen negative at their first exam, or who were unable to access prenatal care earlier, this third trimester screen allows adequate time for persons who seroconvert during their pregnancy to be appropriately treated prior to delivery, and it reduces the risk for congenital syphilis.

**Syphilis screening at delivery:** This requirement is a fail-safe intervention to ensure the syphilis status of newborns is known, and if positive, newborns are provided treatment.

**10. Q: When should syphilis screening occur for a pregnant person if the first examination or the third trimester screening is missed?**

**A:** Pregnant persons who miss their first examination or third-trimester screening should be screened for syphilis as soon as possible.

If a pregnant person has not received prenatal care before delivery, they should be screened at the time of delivery.

Remember, timely syphilis screening is vital to prevent complications and protect the health of both the pregnant person and their baby.

**11. Q: If a pregnant person screens positive for or is diagnosed with syphilis, should they be tested for other sexually transmitted infections, including HIV?**

**A:** Pregnant persons who screen positive for syphilis should be screened for other sexually transmitted infections, including HIV.

**12. Q: Does a pregnant person with syphilis require any special obstetric care?**

**A:** Yes, a pregnant person diagnosed with syphilis will require treatment and follow-up throughout the pregnancy, at the time of birth, and postpartum.

For pregnant persons diagnosed with syphilis in the second half of pregnancy, the Centers for Disease Control and Prevention STI Treatment Guidelines, 2021 recommend the following:

1. A sonographic exam of the fetus. Findings of fetal or placental syphilis should be managed in conjunction with an obstetric specialist.
2. Pregnant patients should be advised to seek care after treatment of syphilis if they experience fever, contractions, or decreased fetal movement due to concern for a Jarisch-Herxheimer reaction, which may be associated with preterm labor and/or fetal distress.

**13. Q: Could a pregnant person be charged with a crime if they are positive for syphilis?**

**A:** No, it is not a crime to acquire syphilis during pregnancy.

## Syphilis Screening

**1. Q: What is considered “standard serological tests for syphilis”?**

**A:** Standard serologic screening for syphilis during pregnancy includes both the traditional (beginning with a quantitative nontreponemal test) and reverse screening algorithms (beginning with a treponemal antibody test). As both screening approaches (traditional and reverse) are valid, neither the Centers for Disease Control and Prevention nor the New York State Department of Health endorses one algorithm over the other.

Per the latest Centers for Disease Control and Prevention Laboratory Recommendations for Syphilis Testing dated February 8, 2024, non-treponemal (lipoidal antigen) and treponemal tests should be interpreted in the same manner regardless of pregnancy status. Practitioners should be aware that in the setting of early primary syphilis (when a chancre first develops), both treponemal and non-treponemal tests may be false negative. In the absence of obvious or visible signs of syphilis and a negative initial treponemal or non-treponemal test, practitioners may perform more frequent testing than the three recommended/required, especially if there are indications of activities that pose a higher risk of syphilis acquisition, such as transactional sex, inconsistent prenatal care, substance use, or having an untested or untreated partner. For additional diagnostic considerations, please see the STI Treatment Guidelines, 2021 from the Centers for Disease Control and Prevention.

**2. Q: Can syphilis point of care testing be used to screen pregnant persons?**

**A:** Yes. Point-of-care tests for syphilis are designed to facilitate quick diagnosis and treatment during a single patient visit. These tests are particularly valuable in settings where traditional laboratory testing may not be readily available or where timely diagnosis is critical. For additional information, please see New York State Guidance [https://www.health.ny.gov/diseases/communicable/std/docs/point\\_of\\_care\\_syphilis\\_test\\_guidance.pdf](https://www.health.ny.gov/diseases/communicable/std/docs/point_of_care_syphilis_test_guidance.pdf)

## Timing of Screening

**1. Q: If a person is diagnosed and adequately treated for syphilis before the third trimester of pregnancy, do they still need to have a third-trimester screen?**

**A:** Yes, the 3rd-trimester syphilis screening requirement is a rule that must be followed.

**2. Q: If the pregnant person has a reactive nontreponemal or treponemal at delivery, but the reactive result is expected and does not relate to untreated syphilis, is it still necessary to screen the newborn?**

**A:** Yes

**3. Q: For scheduled cesarean procedures, does syphilis testing as part of pre-op labs drawn the day prior meet the requirement for syphilis screening at delivery?**

**A:** Yes

**4. Q: If a pregnant person is admitted for delivery, but is discharged home undelivered, does the pregnant person need to be screened again for syphilis when admitted for the actual delivery encounter even if they had a valid third trimester screening?**

**A:** Yes. The third trimester screening is a separate requirement and does not impact the existing requirement to screen at delivery. We suggest working with your organization's general legal counsel regarding the best way to comply with the requirement for screening at delivery, which has not changed.

**5. Q: If a pregnant person delivers before their third trimester, do they still need syphilis screening on admission to delivery?**

**A:** Yes. The delivery screening requirement is still in place.

**6. Q: Should I screen my pregnant patients more than what is required by the laws and regulations?**

**A:** Further screening may be indicated based on the patient risk factors throughout their pregnancy. A centered approach is recommended to promote discussion and share decision making with the patient.

**7. Q: The interim guidance states that pregnant persons “should” be screened between 28 and 32 weeks. Could third -trimester syphilis screening be done before 28 weeks, such as at 24 weeks or between 37 and 38 weeks?**

**A:** The Centers for Disease Control and Prevention recommends screening be done between 28 and 32 weeks. It is advisable for practitioners to use a patient-centered approach to determine the most appropriate timing for third trimester screening that considers the patient’s access to prenatal care and individual risk. This decision-making process may be impacted by other factors such as stigma and health disparities. An open dialogue offers an opportunity to build trust, particularly among marginalized and disenfranchised communities.

**8. Q: If a pregnant person is required to undergo serologic syphilis screening at the time of pregnancy diagnosis, do they have to wait until initiating prenatal care with an obstetric provider for screening?**

**A:** No. Syphilis screening can be performed when the pregnant person presents to an emergency room, clinical or urgent care setting, or other settings where medical care is provided (e.g., correctional facility). New York State recommends screening within those settings to help reduce missed opportunities in preventing congenital syphilis.

**9. Q: If a provider diagnoses pregnancy but is referring the patient for prenatal care, are they required to order the syphilis screen?**

**A:** It is advisable to consult with each health facility’s executive team and/or legal department. Effective communication and integrated care play a crucial role in ensuring that pregnant persons are promptly screened to avoid missed opportunities and prevent health complications. The ideal practice is conduct syphilis screening when the pregnancy is diagnosed. If this is not feasible, a best practice for such providers is to ensure a prenatal appointment, and confirm that syphilis screening and other necessary first trimester pregnancy screenings are scheduled.

**10. Q: If a patient presents for a medication abortion visit in the first trimester and no lab work is being ordered, do they need syphilis screening?**

**A:** No. However, syphilis screening and HIV screening are strongly encouraged, and it is advisable to discuss and support sexual health needs through counseling and follow-up.

## Treatment

**1. Q: What is the recommended treatment for pregnant persons diagnosed with syphilis?**

**A:** Pregnant persons should be treated with the recommended regimen per their specific stage of infection with Penicillin G (Bicillin L-A®). For more information, please refer to the STI Treatment Guidelines, 2021.

*For clinical questions regarding syphilis staging, patient and infant treatment, partner treatment, and recommended clinical follow up for pregnant persons with syphilis in New York State, practitioners/clinical providers can call the Clinical Education Initiative (CEI) line at 1-866-637-2342.*

**2. Q: What if there is a shortage of Bicillin L-A®?**

**A:** If there is a known or anticipated shortage, practitioners should review their existing Bicillin L-A® inventory and reserve Bicillin L-A® for pregnant persons and infants with congenital syphilis. If supply is low, doxycycline is the alternative recommendation that can be used to treat any sexual partner(s) of the pregnant person, and practitioners should closely follow patients to encourage completion of the medication course. However, Doxycycline should **not** be used to treat pregnant persons as it is contraindicated during pregnancy. For guidance and recommendations for syphilis treatment when there is a known shortage of Bicillin L-A, please see New York State Department of Health Advisories: for more information: [https://www.health.ny.gov/diseases/aids/providers/health\\_advisories/](https://www.health.ny.gov/diseases/aids/providers/health_advisories/) and August 2025. Link: [https://www.health.ny.gov/diseases/aids/providers/health\\_advisories/docs/health\\_advisory\\_bicillin\\_recall.pdf](https://www.health.ny.gov/diseases/aids/providers/health_advisories/docs/health_advisory_bicillin_recall.pdf)

## Partner Management

**1. Q: If the pregnant person in my care screens positive for syphilis, what support is there for treating their sexual partner(s)?**

**A:** Partner Services are available, as needed to support the care management of the pregnant persons sex partner(s). For more information about this service, please refer to Accessing Partner Services.

In addition, the New York State Department of Health AIDS Institute funds a robust network of community based agencies that offer screening for sexual transmitted infections. A provider directory is available for finding sexual health and other health related providers. Link: <https://providerdirectory.aidsinstituteny.org/>

**2. Q: Can Expedited Partner Treatment (EPT) be offered to pregnant persons who screen positive for syphilis during pregnancy?**

**A:** The provision of EPT is not permissible for persons diagnosed with syphilis in New York State.

**3. Q: Can doxycycline be prescribed as post exposure prophylaxis to sexual partner(s) of pregnant persons who screen positive for syphilis?**

**A:** Partners of pregnant persons who test positive for syphilis should first be tested and/or presumptively treated for syphilis. After treatment completion, partners may be offered doxycycline (Doxy-PEP) as an STI prevention measure. For more information on Doxy-PEP: Doxycycline Post-Exposure Prophylaxis to Prevent Bacterial Sexually Transmitted Infections (9/25/23) – Clinical Guidelines Program (hivguidelines.org).

## Reporting

### 1. Q: Is it a requirement to report positive syphilis results?

**A:** Yes. For all New York State jurisdictions, physicians are required by law to:

- Report cases to the local health officer (Public Health Law §2101; 10 NYCRR §2.10), and
- Cooperate with state and local health officials' efforts to determine the source and control the spread of sexually transmitted infections (10 NYCRR §2.6).

It is important for the Physician/Offices to report and complete demographic data, especially the pregnancy status of the patient.

### 2. Q: Is there a requirement to report negative syphilis results?

**A: In New York State outside of New York City:** There is currently not a requirement for laboratories to report negative syphilis results.

**In New York City:** An amendment to Health Code § 13.03(b)(2) requires laboratories to report all negative syphilis results. The Health Code previously required the reporting of all positive and indeterminate syphilis results, but only some negative syphilis results. The New York City Board of Health amended the Health Code to enable the collection of all negative syphilis results to improve the diagnosis and treatment of patients and their sexual partners.

## Costs of Syphilis Screening

### 1. Q: Is the cost of these syphilis tests covered by Medicaid?

**A:** For both fee-for-service (FFS) and Medicaid Managed Care (MMC), New York State Medicaid coverage of syphilis screening tests align with syphilis screening requirements during pregnancy. This requires practitioners to order a syphilis screening test for pregnant individuals during their third trimester of pregnancy, in addition to testing at the time of their first exam, and again at delivery.

In addition, for both FFS and MMC, New York State Medicaid will cover point of care syphilis testing in a physician/nurse practitioner/ licensed midwife private office setting. For more information visit: [https://health.ny.gov/health\\_care/medicaid/program/update/2025/no08\\_2025-08.htm#syphilistesting](https://health.ny.gov/health_care/medicaid/program/update/2025/no08_2025-08.htm#syphilistesting)

## Compliance

### 1. Q: What are the Department's expectations around compliance?

**A: In New York State outside of New York City:** With the lack of a reporting requirement for negative syphilis screening results outside of New York City, compliance can only be routinely monitored among pregnancies that result in a congenital syphilis case, which is considered a sentinel event. The New York State Department of Health responds to these sentinel events with timely investigation/review to identify systems, processes, and conditions which caused and/or contributed to the resultant outcome. These investigations/reviews are conducted under the New York State Department of Health's authority pursuant to Public Health Law Article 21 and 10 NYCRR Part 2 as part of public health surveillance, and epidemiologic and patient follow-up. Subsequent recommendations and corrective action, if indicated, may be issued by the New York State Department of Health. Beginning in 2023, all birthing hospitals associated with a reported congenital syphilis case will be issued a notification letter with recommended action items.

Contract entities and other Article 28 facilities might be subjected to retrospective chart reviews to determine compliance.

**In New York City:** The New York City Department of Health and Mental Hygiene responds to reports of congenital syphilis with timely investigation/review to identify systems, processes, and conditions which caused and/or contributed to the resultant outcome. These investigations/reviews are conducted under Article 11 Section §11.03 of the New York City Health Code as part of public health surveillance, and epidemiologic and laboratory investigation activities. Subsequent recommendations and corrective action, if

indicated, may be issued by the New York Department of Health and Mental Hygiene. All health care providers associated with a reported congenital syphilis case are issued a notification letter with recommended action items.

## Information and Resources

### 1. **Q: How can I acquire more information on syphilis screening and treatment, syphilis in pregnant persons, and congenital syphilis?**

- A:**
- For access to free clinical education, or to request training on syphilis or congenital syphilis, please go to [Clinical Education Initiative Training](#). Real-time clinical support is also available through the Clinical Education Initiative Sexual Health Center of Excellence's Clinical line for syphilis and other sexually transmitted infection related questions: Call the **Ask an Expert Clinical Education Initiative Line at 1-866-637-2342**.
  - For access to free clinical materials, including palm cards with information on syphilis in pregnancy and congenital syphilis, and an information sheet on [screening for STIs, HIV, and Hepatitis B/C during pregnancy](#), please go to the [CEI Sexual Health Center of Excellence- Materials Order Form](#).

**New York State:** providers can access information on syphilis and syphilis during pregnancy by visiting the New York State Department of Health [https://www.health.ny.gov/diseases/communicable/congenital\\_syphilis/providers/](https://www.health.ny.gov/diseases/communicable/congenital_syphilis/providers/).

**New York City:** practitioners can access information on syphilis and other sexually transmitted infections and training resources by visiting the New York City Department of Health and Mental Hygiene website: [Sexually Transmitted Infections - Providers - NYC Health](#).

**National:** The Centers for Disease Control and Prevention offers diagnosis and treatment of syphilis and congenital syphilis at [Sexually Transmitted Infections Treatment Guidelines, 2021 \(cdc.gov\)](#). Additional Centers for Disease Control and Prevention resources can be found at [STI Treatment Guidelines \(cdc.gov\)](#) and [STD Facts - Syphilis \(cdc.gov\)](#).



[health.ny.gov](https://health.ny.gov)

