



# Department of Health

KATHY HOCHUL  
Governor

JAMES V. McDONALD, MD, MPH  
Commissioner

JOHANNE E. MORNE, MS  
Executive Deputy Commissioner

**To:** Community Based Organizations, Community Health Centers, County Jails, Emergency Rooms, Substance Use Disorder Treatment Facilities, Urgent Care Centers, Hospitals, Infectious Disease Providers, Internal Medicine Providers, Family Medicine Providers, Local Health Departments, Planned Parenthood, Primary Care Providers, Sexual Health Clinics, and College Health Centers

**From:** New York State (NYS) Department of Health (DOH), AIDS Institute

**Date:** March 10, 2026

Dear Colleagues:

Please see the attached Centers for Disease Control and Prevention (CDC) notification about the availability and use of **Lentocilin**® due to the [shortage of Bicillin® L-A](#) (Penicillin G Benzathine Injectable Suspension).

For those considering purchasing **Lentocilin**®, please use the following options:

- order Lentocillin by setting up a [CostPlus marketplace account](#)
- contact Cost Plus Drug Company at 682-428-8081.

**Lentocilin**® is also a Medicaid reimbursable drug in New York State.

Additionally, we also recommend the use of [DoxyPEP](#) as appropriate in high risk individuals to reduce incidence of new STI infections.

#####



U.S. CENTERS FOR DISEASE  
CONTROL AND PREVENTION

March 10, 2026

Dear Colleagues,

CDC has learned that the U.S. Food and Drug Administration (FDA) is allowing the temporary importation of [Lentocilin®](#) due to the ongoing limited availability [and extended recovery](#) of **Bicillin® L-A** (Penicillin G Benzathine Injectable Suspension, produced by King Pharmaceuticals LLC, a subsidiary of Pfizer). The National Network of STD Prevention Training Centers has posted [information for health care providers on the administration of Lentocilin®](#).

We recommend that jurisdictions strongly consider preserving Benzathine Penicillin G (BPG) for treating patients who are pregnant for several reasons. Penicillin is the only recommended treatment for pregnant women and babies with congenital syphilis. Following the recall and due to high levels of [syphilis diagnoses](#) in the United States, there is currently a very limited supply of BPG.

CDC strongly encourages clinicians and STI programs take the following actions:

1. Take inventory:

- Monitor local supply of BPG and [determine the local pattern of use to forecast need](#).
- Continue to contact distributors to procure BPG as appropriate. [Contact Pfizer to request product](#) for individual pregnant patients or neonates with syphilis through their Medical Request Process if the distributor has no supply.

2. Prioritize using BPG to treat women with syphilis during pregnancy, as penicillin is the only recommended treatment for pregnant women and babies with congenital syphilis.

- Choose doxycycline for men and non-pregnant women to help preserve BPG supplies. Doxycycline can be given at a dose of 100mg orally two times a day for 14 days (for early syphilis) or for 28 days (for late latent syphilis or latent syphilis of unknown duration). See [CDC's treatment guidelines](#) for more information.
- Consider involving antimicrobial stewardship leaders to help institute systems-level approaches to limit the use of BPG and encourage the use of alternative effective antimicrobials for treatment of other infections.
- BPG is one of the preferred prophylactic regimens for patients with acute rheumatic fever or rheumatic heart disease to reduce the likelihood of recurrent Streptococcal infections and acute rheumatic fever episodes. Twice daily administration of oral penicillin V is an alternative option for prophylaxis as detailed in [Clinical Guidance for Acute Rheumatic Fever | Group A Strep | CDC](#).

3. Accurately stage syphilis cases to ensure appropriate use of antimicrobials. Early syphilis (primary, secondary, and early latent) only requires one dose of 2.4 million units of BPG, including during pregnancy. See the [CDC STI Treatment Guidelines](#). A few reminders:

- A thorough physical exam is necessary to accurately stage syphilis. Primary lesions may be hidden inside the mouth, anus, or vagina; signs of secondary syphilis may be found on the skin, mouth, or anogenital

area (i.e., mucus patches, condyloma lata). The rash of secondary syphilis can be subtle.

- Prior syphilis serologic test results can assist with staging a patient with latent syphilis as early latent, if there is evidence of new infection within the last year (i.e., new seroconversion or a sustained four-fold increase in RPR titer in a person who has had syphilis in the past).
- Reviewing signs and symptoms and sexual partner history from the past 12 months can assist with determining the likelihood of syphilis acquisition within the last 12 months, which also only requires one dose of Bicillin® L-A (2.4 million units) instead of three doses (7.2 million units).
- You can find resources for appropriately staging syphilis from [the National STD Curriculum](#) (NSTDC) and [the New York City Department of Health and Mental Hygiene Bureau of Sexually Transmitted Infections/New York City STD Prevention Training Center](#).

4. Communicate with healthcare providers and pharmacists.

- Notify healthcare providers and pharmacists of the limited availability of BPG so they are aware and can plan appropriately.
- Clinicians with questions about syphilis clinical management should contact an Infectious Diseases specialist or [the online National Network of STD Clinical Prevention Training Centers \(NNPTC\) STD Clinical Consultation Network](#).

5. Notify Division of STD Prevention and FDA of any shortage of BPG so we can continue to monitor this situation.

- If you are a *Strengthening STD Prevention and Control for Health Departments* (STD-PCHD) funding recipient and have questions about any of this information or need assistance, please contact your project officer.
- For other groups who are encountering challenges securing this drug to treat women with syphilis during pregnancy, please contact [stdshortages@cdc.gov](mailto:stdshortages@cdc.gov).

We appreciate all of your hard work to prevent and address syphilis.

Sincerely,

**Lindley Barbee, MD, MPH**

Chief Medical Officer (Acting)  
Division of STD Prevention  
National Center for HIV, Viral Hepatitis, STD and TB Prevention  
Centers for Disease Control and Prevention  
U.S. Department of Health and Human Services

**Puja Seth, PhD**

Director (Acting)/ Principal Deputy Director  
Division of STD Prevention  
National Center for HIV, Viral Hepatitis, STD and TB Prevention  
Centers for Disease Control and Prevention  
U.S. Department of Health and Human Services

Share this link to invite colleagues to subscribe to receive HIV, Viral Hepatitis, STI, and TB News & Updates | CDC and follow the two-step email verification process.