

Levels of care for maternal and neonatal healthcare: a scoping review

Shaina Mackin^{1*}, Louise Tina Day^{2*}, Caity Dekker¹, Isabel Gouse¹, Michuki Maina³, Jalemba Aluvaala^{4†}, Allisyn Moran^{5†}; Revisioning Emergency Obstetric and Newborn Care (EmONC) levels-of-care workstream

¹Averting Maternal Death and Disability, Columbia University Mailman School of Public Health, New York, New York, USA

²Department of Infectious Disease Epidemiology and International Health, London School of Hygiene & Tropical Medicine, UK

³KEMRI Wellcome Trust Research Programme, Nairobi, Kenya

⁴KEMRI Wellcome Trust Research Programme and Department of Paediatrics and Child Health, University of Nairobi, Nairobi, Kenya

⁵Sexual, Reproductive, Maternal, Child, Adolescent and Ageing Health Department, World Health Organization, Geneva, Switzerland

*Joint first authorship.

†Joint senior authorship.

Background To meet the 2030 aims of the Global Strategy for Women's, Children's and Adolescents' Health and Sustainable Development Goals, annual rates of maternal and newborn mortality and stillbirth must decrease. The organisation of maternal and newborn health (MNH) services influences access to and quality of care. We designed this scoping review to understand how levels of MNH care are organised in different country contexts.

Methods We conducted a scoping review of peer-reviewed literature published after 2009. Based on the World Health Organization (WHO) quality-of-care framework, we conducted descriptive and deductive textual narrative analysis to identify the reported number of levels of MNH care stratified by country and mortality rates; describe how levels are conceptualised; and explore alignment of levels for the maternal-newborn dyad.

Results We included 162 of 3591 reports. The number of MNH facility levels of care across 56 countries ranged from two to seven. Types of identified MNH care facilities were described at varying levels. Two levels of care were reported in 5% of cases, three levels in 55%, four levels in 30%, five levels in 8%, and six and seven levels in <1% of cases. Home and community-based MNH care (non-facility) was reported in 8% of country descriptions. Countries with the lowest stillbirth, maternal, and newborn mortality rates mostly reported three or four facility levels. The criteria used to distinguish MNH levels of care as low, intermediate, and high were aligned with domains of the WHO quality-of-care framework, mostly human and physical resources.

Conclusions Levels of MNH care described in the literature were distinguished by characteristics, including provision and experience of routine and emergency care. Three or four levels of MNH facility care were most commonly reported. Linking maternal and newborn facility care to community connections is an important consideration to ensure equitable access to routine and emergency care.

Correspondence to:

Louise T Day
London School of Hygiene and Tropical Medicine (LSHTM), Infectious Disease Epidemiology and International Health
Keppel Street, London
UK
louise-tina.day@lshtm.ac.uk

Shaina Mackin
Averting Maternal Death and Disability (AMDD), Columbia University Mailman School of Public Health
722 West 168th Street, New York
USA
shaina.mackin@gmail.com

Increasing access to institutional childbirth has been a major strategy for several decades to end preventable maternal and newborn deaths and stillbirths. Although >80% of global births currently occur in health facilities [1], mortality rates are decreasing too slowly to meet the Global Strategy for Women's, Children's and Adolescents' Health and the Sustainable Development Goals [2].

The organisation of the maternal and newborn health (MNH) care affects women's and newborns' access to care, and currently, access to MNH services remains inequitable worldwide. Emergency clinical interventions are necessary to prevent death for the most common direct causes of death and disability for both women (e.g. haemorrhage, eclampsia, and sepsis) and newborns (e.g. intrapartum-related complications, sepsis, and preterm birth) [3–5]. The



emergency obstetric care (EmOC) framework, used since 1997 in high-mortality/morbidity settings to plan the availability and accessibility of emergency interventions, is currently being revised [6–9]. The existing framework conceptualises two levels of EmOC health facilities based on the reported performance of tracer ‘signal functions,’ or emergency clinical interventions, within a specified timeframe. These two levels are comprehensive EmOC (CEmOC), in which all nine signal functions are performed, and basic EmOC (BEmOC), including a subset of seven signal functions [8]. In 2012, researchers proposed measuring a set of newborn-specific ‘signal functions’ across different levels of emergency care [10]. In 2018, the World Health Organization (WHO) and United Nations Children’s Fund (UNICEF) proposed levels of care for small and sick newborns, also based on clinical intervention availability [11].

The organisation of MNH care is essential to delivering high-quality services, and the poor quality of care is a major contributor to preventable deaths of women, newborns, and stillbirths [12]. The 2015 WHO vision of quality of care for MNH articulates the structures and processes needed for people-centred health outcomes, including the provision and experience of care [13,14]. The resulting framework articulates eight interconnected domains: evidence-based practices, actionable information systems, functional referral systems, effective communication, respect and dignity, emotional support, competent and motivated human resources, and essential physical resources.

Health service delivery is typically organised into levels that vary by setting – primary, secondary, and tertiary. Improving MNH care services organisation is an important lever for change to accelerate progress in ending preventable mortality and morbidity among women and children. As illustrated by the launch of the ‘Every Woman Every Newborn Everywhere’ platform, this includes a focus on aligning and integrating maternal and newborn service delivery to end preventable mortality and morbidity [15]. Recent WHO guidance on networks of care for MNH describes the importance of connections between levels of MNH care [16]. However, little is known about the optimal levels of care in the health system for high-quality MNH care, both for routine care and for care when complications arise.

In this research, we aimed to explore how countries, since the 2009 EmOC revision, organised MNH care into levels. We collaboratively designed this study with the project Revisioning EmONC ‘Levels of Care’ workstream’s technical advisors [9]. We had three goals: to report how many levels of MNH care have been organised in different country health systems and stratify them by mortality rates, to describe the criteria used to define and conceptualise maternal and newborn levels of care within country health systems, and to explore alignment of levels of care for the maternal-newborn dyad (Table 1).

Table 1. Objectives and research questions

Objectives and research questions
To report how many levels of MNH care are operationalised in country health systems
<i>How many levels of MNH care are operating in different countries’ health systems?</i>
To describe the criteria used to define and conceptualise MNH levels of care within country health systems
<i>How have experts defined levels of MNH care?</i>
<i>How have health systems organised levels of MNH care?</i>
<i>How does regionalisation fit into the definition and organisation of levels of MNH care?</i>
<i>How do referrals fit into the definition and organisation of levels of MNH care?</i>
<i>Can the domains of the WHO’s framework for the quality of MNH care be applied to defining and organising levels of MNH care?</i>
To explore the alignment of levels of care for the maternal-newborn dyad
<i>How have health systems and experts conceptualised the maternal-newborn dyad in the context of levels of care?</i>
To list measures identified to determine levels of MNH care
<i>What indicators do health systems and experts use for maternal and newborn levels of care?</i>

MNH – maternal and newborn health, WHO – World Health Organization

METHODS

We designed this scoping review using the updated Arksey and O'Malley framework and reported its findings in accordance with the PRISMA-ScR [17–21] (Checklist S1 in the [Online Supplementary Document](#)).

Search strategy and eligibility criteria

We reviewed peer-reviewed publications across all income settings to provide broad evidence on maternal and newborn levels of care. In October 2021, we searched four bibliographic databases – MEDLINE, Embase, Cochrane, and Google Scholar – using a search strategy designed collaboratively by the Revisioning EmONC project's working group (Table S1 in the [Online Supplementary Document](#)). For feasibility, we limited Google Scholar results to the top 200 results. To ensure this did not lead to the omission of relevant literature, at the conclusion of title and abstract screening, from October 2021 to August 2022, we sent the list of included records to the Steering Committee of the Revisioning EmONC project and committee members to solicit feedback and identify potential studies that may have been omitted, based on their subject matter expertise. English, Spanish, Portuguese, French, and Arabic were included as eligible languages.

We used two search strings across all databases. Eligibility for the first string, which included MNH and levels of care keywords, was restricted to records published from 2009 onwards to reflect developments in the MNH levels of care literature since the last revision of the EmONC handbook in 2009 [8]. The second string combined two search terms from the 2020 WHO, United Nations Population Fund, and UNICEF review of maternal networks of care [22]. This search was restricted to publications after June 2020, when the landscape review was published, to identify updates in the literature (Table S2 in the [Online Supplementary Document](#)).

Data management, extraction, and analysis

After identifying and removing duplicates, two primary reviewers (CD and SM) double-screened the remaining titles and abstracts. They initially calibrated their screening decisions by comparing and reaching consensus on a subset of records to ensure consistent application of the inclusion criteria and use of the data extraction tool (Table S3 in the [Online Supplementary Document](#)). Conflicts over whether records should be included or excluded were resolved by a third reviewer (LD). An additional reviewer with the necessary language proficiency (IG) screened Spanish records and then discussed with the primary reviewers (SM and CD) to decide on inclusion or exclusion. CD and SM conducted full text review and dual data extraction using a customised extraction tool in Covidence (Covidence, Melbourne, Victoria, Australia) for the following variables: study design, type of literature, units of analysis, client focus (*i.e.* maternal, neonatal), geography by WHO regions, number of MNH levels of care, descriptions of levels of care, discussion of the maternal-newborn dyad, and measures used to determine levels of care. Possible conflicts between reviewers (CD and SM) were resolved in consultation with a third researcher (LD).

We imported publicly available WHO/UNICEF data on maternal and neonatal mortality and stillbirths by country from the Healthy Newborn Network and merged them with extracted data to map MNH outcomes to the number of levels of care [23]. We then summarised, by outcomes, the number of MNH care levels operating in country settings. We categorised reports that described levels of care in country settings but did not explicitly state or describe how many levels existed as having an 'unspecified' number of levels.

We analysed qualitative data using textual narrative synthesis, conceptualising the criteria used for defining MNH levels of care. We primarily used a deductive analysis, with the domains of the WHO quality-of-care framework for MNH (modelled according to the Donabedian flow of structure, process, and outcome) as *a priori* nodes. We began with a codebook consisting of the framework's domains. Following consensus among CD, SM, and LD, additional codes were introduced during textual narrative analysis as themes emerged from the literature. All codes mapped to the WHO framework for the quality of MNH care are presented in [Figure 1](#) [4,24,25]. Moreover,




Figure 1. Results from textual narrative synthesis mapped to WHO framework for the quality of maternal and newborn health care [14].

we generated descriptive summary statistics. The findings were discussed and reviewed by the Revisioning EmONC levels of care technical working group and Steering Committee to validate the relevance of the synthesised findings to strengthen credibility (Table S1 in the [Online Supplementary Document](#)).

We managed data using Zotero (Corporation for Digital Scholarship, Falls Church, Virginia, USA) reference manager for search results, Covidence (Covidence, Melbourne, Victoria, Australia) software for workflow management, R, version 4.1.1 (R Core Team, Vienna, Austria) for quantitative analysis, and NVivo, version 12 (Lumivero, Denver, Colorado, USA) for qualitative data.




Figure 2. PRISMA flowchart of included reports. *For feasibility purposes, Google Scholar results were limited to the top 200 results.

tals, etc.). Quantifying the number of facility-based levels (excluding home and community-based 'non-facility' levels reported by studies), two, three, four, five, six, and seven levels of care were described in 8 (5.3%), 83 (54.97%), 46 (30.46%), 12 (7.95%), and 1 (<1%) of cases, respectively (Figure S1 and Table S5 in the [Online Supplementary Document](#)).

Different reports sometimes conflicted on the number of MNH levels of care described in the given countries. In ten out of 56 (17.9%) countries (Australia, Canada, Ghana, Nepal, South Africa, Spain, Tanzania, Turkey, the UK, and the USA), different reports reported different numbers of MNH levels. For example, Tanzania was said to have five levels by some reports, and three by others.

RESULTS

Selection of sources and summary statistics

We retained 389 of 2499 records for full-text review after title and abstract and retained 162 for data extraction ([Figure 2](#); Table S4 in the [Online Supplementary Document](#)).

More than half of the included reports were from the WHO Regions of the Americas (56%) and Africa (32%), with 89% of reports from the Americas reporting on the USA or Canada ([Figure 3](#)). The studies predominantly had observational (65%) and secondary analysis (33%) designs. The unit of analysis varied from multinational regions to countries to health facilities.

Objective one: identifying the number of MNH levels of care and stratifying by mortality rates

Based on 121 (74.7%) of the 162 included reports, the number of MNH levels of care ranged between two and seven, with a median of four. The remaining 41 reports (25.3%) discussed concepts related to MNH levels of care without specifying a number. Reported MNH levels of care included 'home' (with or without provider presence), 'community' (unspecified whether at home or in basic facility), and 'health facility' levels (e.g. birthing centres, health posts, primary care facilities, hospitals, etc.).




Figure 3. Characteristics of included reports regarding levels of care for maternal and neonatal health-care (n=162). The map presents the number of reviews per country.

Most of the lowest maternal, stillbirth, and neonatal mortality settings, including Denmark, Norway, Poland, Spain, and Italy, were described as having three levels of care (Figure 4; Table S6 in the Online Supplementary Document) [23].

Objective two: defining and conceptualising MNH levels of care

Through the thematic analysis, we identified MNH levels of care as the lowest, intermediate, and highest levels of the health system. These levels had varied conceptualisations when mapped onto the structure, process, and outcome components of the WHO quality-of-care framework for maternal and newborn healthcare (Figure 1, Table 2) [14]. Some reports explicitly used the label 'level 1' for the lowest level of care and 'level 2', 'level 3', *etc.* for higher levels.




Figure 4. Number of MNH facility levels of care identified by outcomes.

Table 2. Number of codes identified through thematic analysis mapped to the WHO quality-of-care framework among included papers (n = 162)

Codes	Occurrences	Papers
Health systems	18	13
Maternal-newborn dyad	56	31
Highest level of care	70	54
Lowest level of care	83	57
Domain 1: evidence-based practices	3	3
Clinical patient characteristics	110	71
EmOC	67	28
Risk-appropriate care	58	39
Domain 2: actionable information systems	4	3
Telemedicine	24	10
Domain 3: functional referral systems	23	17
Distance or geography	5	5
Referral	103	51
Regionalisation	162	78
Transport	75	39
Domain 4: effective communication	29	18
Domain 5: respect and preservation of dignity	8	7
Domain 6: emotional support	3	3
Domain 7: competent, motivated human resources	31	24
HR characteristics	110	72
Domain 8: essential physical resources available	21	17
Facility characteristics	157	83
Outcomes	64	38
Demographic patient characteristics	2	2
Volume	51	33

EmOC – emergency obstetric care, HR – human resources

ers to specify a high-level health facility designation (e.g. tertiary) [35,37]. Occasionally, reports cited countries that consider primary health centres to be the highest level [38]. Further, 54 papers (33.3%) discussed ‘highest levels of care,’ while 57 papers (35.2%) discussed ‘lowest levels of care.’

Process

We identified descriptors used to demarcate levels of care mapped to all the quality domains and dimensions of the quality-of-care framework [13,14].

Provision of care

Evidence-based care (quality domain one) was used to delineate low, intermediate, and high categories of MNH services based on clinical conditions or risk-stratification [39]. Highest category

Structure

The level of MNH care was frequently categorised as low, intermediate, or high, though it was defined differently across settings. Most reports considered the lowest level (level 1) to be ‘health facilities’, typically falling below full hospitals and including primary health centres and community-based services. Examples included: community-based health planning and services and/or maternity homes (e.g. Ghana) [26–28], dispensaries (e.g. Tanzania) [29], community hospitals and/or birthing centres (e.g. USA and Canada) [30–34], and sub-base hospitals (e.g. Japan) [35].

From the included reports, 17 (10.5%) referenced non-facility (e.g. home or community) levels of MNH care at the lowest level of their health systems. In some of these cases, a three-level service delivery model was depicted: family and community (low level), outpatients and outreach clinics (intermediate level), and health facilities and hospitals (high level) [35]. Some reported that MNH care was structured as either home births or ‘level 1 facilities’, being considered the lowest level of care (e.g. UK) [36].

The category ‘high’ also varied in application across places of care, with some health systems using it to refer to any health facility or hospital, and oth-

services were conceptualised as care bundles for women and newborns with complications (e.g. blood transfusion and caesarean section for CEmOC) [40–44]. Some reports stratified the lowest level of care offered within their highest facility category as such (e.g. a ‘level I neonatal intensive care unit’) within an intensive care unit that also contains levels II–IV neonatal intensive care unit care) [45–51]. Further, 71 papers (43.8%) discussed evidence-based care in terms of ‘clinical patient characteristics.’

Lower-level categories of care referred to a range of service provision, including childbirth services with essential newborn care. One report from Kenya noted that the lowest levels may be the most neglected [39]. Another report describing Nepal’s health system stated that only the highest-level facilities have the capacity to offer even basic EmONC care [52]. One report proposed that established levels of intensive care be adapted for maternal care, with routine maternity care services at the lowest level [53].

Actionable information systems (quality domain two) were described in the reports in terms of the process linking levels of maternal and newborn care. Telemedicine emerged as a means of creating networks to improve the functioning of care levels. Examples were given from the start of the COVID-19 pandemic, showing how new telemedicine interfacility intensive care specialist consultation was used to overcome physical barriers to care during lockdowns, providing higher levels of intensive or specialised, family-centred care [45]. Actionable information systems were also described as contributing to referral efficiency between sending and receiving facilities across levels of MNH facility care [54–56]. No standardised indicators determining maternal and/or neonatal levels of care were found among the identified reports that distinguished between MNH levels of care.

Functional referral systems (quality domain three) also provide the critical linkage between maternal and newborn levels of care. Reports frequently used related descriptors at the hospital and health facility levels, including geography, referral, transfer, transport, travel time, distance, and ‘regionalisation’ (e.g. regional hospital or centre). A recurrent theme was the importance of strengthening referral linkages, including coupling preventive community work with improving the quality of facility care, as well as strengthening transport linkages before, during, and after facility births [57,58]. Further, we identified 51 papers (31.5%) discussing ‘referral’ and 78 papers (48.1%), ‘regionalisation.’

Experience of care

Some descriptors were found, though fewer than in other domains, relating to levels of care that included the dimension of experience of care through respect and preservation of dignity [13,14].

Effective communication (quality domain four) was not frequently reported. One Canadian study discussed the importance of provider-family communication during transfers between levels of care [59]. Otherwise, effective communication was only discussed in terms of its absence being a bottleneck to effective MNH care [60,61].

Respect and preservation of dignity (quality domain five) in levels of MNH care was established through discussion of respectful maternity care, with included papers also citing ‘family-centred’ [45], ‘patient-centred’ [62], and ‘client-centred’ [63] care. Separation of mother and baby due to one of the two requiring a higher level of care was reported to cause feelings of frustration and disrespect [64].

Emotional support (quality domain six) emerged as a component of holistic maternal care in relation to the social and psychological impacts of pregnant women categorised as obstetric high-risk being transferred from their community to a higher level of care [65]. In relation to emotional support, one report specifically designated four levels of evidence-based psychological care for women experiencing obstetric complications based on their conditions [66].

Additional quality processes

Competent and motivated human resources (quality domain seven) were frequent descriptors for MNH levels of care, mentioned in 72 papers (44.4%).

One report from a conflict-affected area defined the highest level of care not by place or intervention, but by the skill level of the human resources (*e.g.* maternal health workers) [67]. Other reports linked the health worker cadre with the place of care to describe the lowest level of care, predominantly in higher mortality settings: Traditional birth attendants at health huts or health posts [67,68], skilled birth attendants at home or in community health centres [43], midwives at midwife obstetric units [60,63], community health workers running community-based health planning and services zones [26–28], and ‘village health team/services’ or female community health volunteers operating in different health systems [69]. There was high variation between countries in the cadre at the lowest level of MNH care.

Essential physical resources (quality domain eight) were also frequent descriptors, for level-specific MNH facility care, found in 83 papers (51.2%). Collectively, physical resources distinguished newborn ‘special care’ from ‘neonatal intensive care units.’ Specific examples included routine ward care for low-risk patients [70] and wards equipped with extracorporeal membrane oxygenation equipment for the sickest patients [71].

Outcomes

We found that person-centred and facility outcomes were related in various ways to levels of MNH care in the reports identified. Patient volume was commonly described in terms of outcomes (*e.g.* obstetric haemorrhage), and the levels of care hierarchy were described in terms of facility volume and output [72]. Facility-level drivers for improved outcomes included strengthening lower levels of care to reduce neonatal mortality rates [73] and increasing in-utero referral and transport between levels of care to ensure better outcomes than postpartum referral and transport [50]. Patient demographic characteristics, a salient theme *in vivo*, were frequently associated with MNH outcome stratification. One report from the USA identified neglecting to escalate clients of colour to risk-appropriate levels of care as a determinant of disparities in MNH outcomes [74]. Three predominant categories of characteristics were used to distinguish between hierarchical levels of maternal and neonatal care across included reports: facility characteristics (cited 157 times across 83 papers), clinical patient characteristics (cited 110 times across 71 papers), and human resource characteristics (cited 110 times across 72 papers). Volume and geography/distance were also cited in distinguishing levels, though they are not as salient [50]. Similarly, potential bottlenecks in the implementation of best emergency obstetric care practices included a lack of personnel or competence among them, inadequate essential physical resources (*e.g.* products and/or technologies), and high patient volume [43,54].

Objective three: exploring levels of care for the maternal newborn dyad

Among 162 reports, levels of care were described in 63 (38.9%) for neonatal care, 50 (30.9%) for maternal care, and 44 (27.2%) reports for both maternal and neonatal care. The remaining five reports (3.1%) discussed concepts related but not explicit to MNH levels of care (*e.g.* regionalisation of care, networks of care, *etc.*).

Several papers highlighted the concept of the maternal-newborn dyad and how the mother and neonate move together or separately through levels of care [34,75–77]. In general, correlation between levels of care for women and newborns was advocated, whilst acknowledging the challenges of correlating variation in criteria to distinguish levels and regionalisation of care for neonatal *vs.* maternal care [78–80]. Several reports from higher-income settings state levels and regionalisation for neonatal care have received more attention than obstetric care [31,79,81,82]. The aspiration to integrate a maternal-neonatal network or system of levels of care was a stated priority to ensure optimal dyad outcomes [31,33,38,83] and to avoid fragmentation of care and

resources [47]. Extra emphasis on *in utero* transfer was sometimes articulated, as was prioritisation of the woman or baby for transfer to a higher-level facility [82]. Very few reports argued against the designation of levels of care coordinated between mother and foetus [31,84] to prioritise the mother [33]. Mostly, examples were given of efforts to coordinate more effectively (e.g. USA has recently made efforts to incorporate maternal care into neonatal levels of care) [34,76,77].

DISCUSSION

Summary of evidence

In this scoping review, we summarised peer-reviewed literature regarding MNH levels of care (n=162) published between 2009 and 2021. We found three to four levels of facility MNH care the most commonly reported, including those from settings with the lowest stillbirth, maternal, and neonatal mortality rates. In these reports, levels of care were characterised by both routine and emergency care criteria, across many domains of the WHO quality-of-care framework [13,14]. These findings can be used to strengthen the health system and accelerate progress for women and newborns towards the globally agreed goals.

Reported MNH levels of care varied in both the number reported within and between countries. These discrepancies could be due to health system reforms, parallel ministries or systems, differing definitions of levels of care, descriptions including *vs.* excluding non-facility levels of care, descriptions of emergency-specific designations (e.g. BEmONC/CEmONC) *vs.* MNH care, including routine and emergency services, maternal/neonatal care being described together *vs.* separately, or descriptions of recommended *vs.* actual operationalised levels of care (Table S7 in the [Online Supplementary Document](#)).

The lowest level of MNH care varied by context, with some contexts including non-facility levels and others excluding them. The centrality of MNH care within community healthcare was evident in the identified literature [85]. Emphasis on improving access to emergency care and institutional deliveries should not preclude recognition of the importance of community-based care. Ensuring high-quality care across the ‘household to hospital continuum of care’ is critical to achieve universal coverage by integrating primary healthcare with higher-level facility care [86].

Levels of MNH care described in these manuscripts included both routine and complication-related care. For emergency-specific designated levels of care, the two-level BEmONC and CEmONC structure was mentioned in only six reports since 2009. The EmOC framework was designed in 1997, reaffirmed in 2009, and is currently being revised [7-9], with the two levels of care determined by tracer clinical emergency intervention signal functions. The highest level, CEmOC, is achieved by performing caesarean sections and blood transfusions, in addition to the BEmONC signal functions [9]. Specifically, critical and/or intensive care has emerged as an empirical, categorical need across MNH systems to care for the sickest women and newborns [39,42]. As countries transition across the maternal newborn stillbirth transition, further reductions in preventable maternal and newborn mortality and stillbirth rates will not be possible without access to high-quality special and intensive care [87].

Approaching MNH levels of care while considering quality of care, in both experience and provision, is important for improved outcomes for women and newborns. We found that countries’ definition of their MNH levels of care was articulated with other characteristics, beyond tracer clinical emergency interventions. These characteristics aligned with the WHO quality-of-care framework, including human and physical resources [13,14]. Recognising that poor quality of care is a major contributor to preventable mortality and morbidity, measuring other criteria beyond signal functions may contribute to the improvement in quality of care needed. Multidimensional quality of care is needed at every level of the health system, and describing level-specific quality domains could be helpful for programme planning.

High-quality MNH care is safe, effective, timely, efficient, equitable and people-centred [13]. We found that integrating maternal and newborn people-centred care across MNH levels of care, emphasising the importance of avoiding separation of the mother and baby, leads to better maternal and newborn outcomes. Historically, the individual woman or newborn has been prioritised in regionalisation efforts, with neonatal care levels receiving more attention than maternal care levels [31,79,81,82]. Instead, a contextualised, networked maternal-neonatal dyad-levels-of-care approach is needed [64,78]. This aligns with the recent focus on strengthening respectful and family-centred care. Integrated MNH levels of care should be included in emerging conceptualisations of health system organisation, using a programmatic transition framework [87].

Strengths and limitations

To our knowledge, our study is the first scoping review specific to levels of maternal and neonatal care; however, it has several potential limitations. We introduced potential language bias by restricting publications to English, Spanish, Portuguese, French, and Arabic, which could bias results toward regions where these languages are predominantly spoken. Google Scholar search results were limited to the first 200 due to the high volume. We focused on peer-reviewed publications to identify the latest evidence on levels of care thinking, but we recognise this might have excluded relevant sources and contributed to publication bias. It was beyond the scope of this study to review grey literature, including country MNH and wider health system policy documents, to understand how MNH levels of care are incorporated within the health system. We recognise that excluding grey literature may have led to the omission of valuable data, such as government reports and policy documents, which can offer insights that peer-reviewed studies alone may not.

Although the study had a global focus, the identified studies were not balanced across all geographies, with more than 50% from the Americas (and among these, many from North America), 30% from Africa, and fewer from Europe and Asia. The publications we identified spanned a 12-year period, and the country mortality rates we used to stratify may reflect more recent revisions to MNH levels of care. In line with scoping review methodology, we did not formally assess the quality of the evidence included, as is typically done in systematic reviews.

In addition, robustly addressing the role of inequity for marginalised groups and structural inequities was outside the scope of this review. The literature reviewed in this study lacked coverage of racial and ethnic disparities in MNH care, and the authors strongly feel that more research is needed on this topic as it relates to access to the levels of care available. For example, race and ethnicity are vital characteristics in the country-specific operationalisation of levels of care. Further research into equitable risk-appropriate escalation to appropriate MNH facility levels of care is paramount to the advancement of reproductive justice across maternal and neonatal outcomes.

CONCLUSIONS

In scoping the literature on MNH levels of care, there is variation within and between countries in the number of levels described and the ways they are characterised. Among the identified countries, three or four levels of care were most often reported, including those in the lowest mortality settings. Levels of MNH care are characterised by multiple domains of the WHO quality-of-care framework, including human and physical resources, interventions, and referral structures from the community to the facility. While the dimensions of MNH care provision may differ by level, the experience of care for women and newborns needs to be prioritised at every level of care. Integration of care for the maternal-newborn dyad is a needed focus which will support family-centred care with better outcomes for all. As policymakers and practitioners plan MNH care services, these results may inform health system organisation across three or more levels of care, including intensive care at the highest level. Integration of maternal and newborn services must be a priority across all geographies to strengthen person- and family-centred care and accelerate progress for every woman and every newborn everywhere.

Revisioning EmONC Levels of Care workstream: Allisyn Moran, Jalemba Aluvaala, Louise Tina Day, Shaina Mackin, Caity Dekker, Michuki Maina, Isabel Gouse, Patience Afulani, Patsy Bailey, Lynn Freedman, Tedbabe Hailegebriel, Elimase Kamanga, Mary Kinney, Ornella Lincetto, Samantha Lobis, Zoe Matthews, Jean-Pierre Monet, Isabelle Moreira, Sarah Moxon, Kristine Nilsen, Cynthia Boschi, Kate Ramsey, Jennifer Requejo, Sanam Roder-DeWan, Suzanne Stalls, Caitlin Warthin. The full list with affiliation is available in the [Online Supplementary Document](#).

Acknowledgements: We thank Kaosar Afsana for her contributions to the Revisioning EmONC Levels of Care workstream.

Disclaimer: The authors have been given permission to publish this article. The authors alone are responsible for the views expressed in this publication, and they do not necessarily represent the views, decisions, or policies of the WHO.

Ethics statement: This study was a review of existing literature; thus, ethical approval was not sought.

Funding: This work was supported, in whole or in part, by the Bill and Melinda Gates Foundation (INV-001363). The funders (the Foundation and USAID) did not have a role in the study design, data collection, management, analysis, interpretation, writing of the manuscript, or the decision to submit manuscripts for publication. Under the grant conditions of the Foundation, a Creative Commons Attribution 4.0 generic license has already been assigned to the authors' accepted manuscript version that might arise from this submission. The article processing charge was funded by the WHO Sustained Growth Fund.

Authorship contributions: Conceptualisation: ACM and JA, in consultation with the Revisioning EmONC Levels of Care Working Group. Data curation: SM, CD, and LTD. Formal analysis: SM and LTD. Methodology: ACM, JA, LTD, and SM. Project administration: ACM. Supervision: ACM and JA. Writing – original draft: SM and LTD. Writing – review and editing: SM, LTD, CD, IG, MM, JA, AM, Revisioning Emergency Obstetric and Newborn Care (EmONC) Levels of Care workstream.

Disclosure of interest: The authors completed the ICMJE Disclosure of Interest Form (available upon request from the corresponding author) and declare the following activities and relationships: Shaina Mackin, Caity Dekker, Isabel Gouse, research assistant fees and Michuki Maina, Jalemba Aluvaala consulting fees from Averting Maternal Deaths and Disabilities from Averting Maternal Deaths and Disabilities. Louise T Day grant from the WHO. Allisyn Moran is employed by the WHO and had funding from the Bill and Melinda Gates Foundation.

Additional material

[Online Supplementary Document](#)

REFERENCES

- 1 United Nations Children's Fund. Delivery Care. 2024. Available: <https://data.unicef.org/topic/maternal-health/delivery-care/>. Accessed: 20 January 2026.
- 2 World Health Organization. Improving maternal and newborn health and survival and reducing stillbirth: progress report 2023. Geneva, Switzerland: World Health Organization; 9 May 2023. Available: <https://www.who.int/publications/i/item/9789240073678>. Accessed: 20 January 2026.
- 3 World Health Organization, United Nations Children's Fund, United Nations Population Fund, World Bank Group, UNDESA/Population Division. Trends in maternal mortality 2000 to 2020: estimates by WHO, United Nations Children's Fund, UNFPA, World Bank Group and UNDESA/Population Division. Geneva, Switzerland: World Health Organization; 7 April 2025. Available: <https://www.who.int/publications/i/item/9789240108462>. Accessed: 20 January 2026.
- 4 United Nations Inter-agency Group for Child Mortality Estimation (UN IGME). Never Forgotten: The situation of stillbirth around the globe. New York, New York, USA: United Nations Children's Fund; 9 January 2023. Available: <https://data.unicef.org/resources/never-forgotten-stillbirth-estimates-report/>. Accessed: 20 January 2026.
- 5 United Nations Inter-agency Group for Child Mortality Estimation (UN IGME). Levels & Trends in Child Mortality: Report 2022, Estimates developed by the United Nations Inter-agency Group for Child Mortality Estimation. New York, New York, USA: United Nations Children's Fund; 2022. Available: https://www.un.org/development/desa/pd/sites/www.un.org.development.desa.pd/files/undesa_pd_2022_un-igme_child_mortality_report.pdf. Accessed: 20 January 2026.
- 6 Maine D, McCarthy J, Ward V. Guidelines for monitoring progress in the reduction of maternal mortality. New York, New York, USA: UNICEF; 1993. Available: <https://www.publichealth.columbia.edu/file/10730/download?token=LzW5VePB>. Accessed: 10 March 2026.
- 7 Maine D, Wardlaw TM, Ward VM, McCarthy J, Birnbaum A, Akalin MZ, et al. Guidelines for monitoring progress in the reduction of maternal mortality. New York, New York, USA: UNICEF; 1997. Available: <https://www.publichealth.columbia.edu/file/10730/download?token=LzW5VePB>. Accessed: 10 March 2026.

- 8 World Health Organization, United Nations Population Fund, United Nations Children's Fund, AMDD. Monitoring emergency obstetric care: a handbook. Geneva, Switzerland: World Health Organization; 2009. Available: <https://www.who.int/publications/i/item/9789241547734>. Accessed: 20 January 2026.
- 9 Averting Maternal Death and Disability. Re-visioning Emergency Obstetric and Newborn Care Project Brief. New York, New York, USA: Columbia University; 2020. Available: <https://emonc.org/assets/tools/en/re-visioning-evidence-base.pdf>. Accessed: 20 January 2026.
- 10 Gabrysch S, Civitelli G, Edmond KM, Mathai M, Ali M, Bhutta ZA, et al. New signal functions to measure the ability of health facilities to provide routine and emergency newborn care. *PLoS Med*. 2012;9:e1001340. **Medline:23152724 doi:10.1371/journal.pmed.1001340**
- 11 World Health Organization, United Nations Children's Fund. Survive and thrive: transforming care for every small and sick newborn. Key findings. 2018. Available: <https://www.unicef.org/reports/transforming-care-for-every-small-and-sick-newborn-2020>. Accessed: 20 January 2026.
- 12 Kruk ME, Gage AD, Arsenault C, Jordan K, Leslie HH, Roder-DeWan S, et al. High-quality health systems in the Sustainable Development Goals era: time for a revolution. *Lancet Glob Health*. 2018;6:e1196–252. **Medline:30196093 doi:10.1016/S2214-109X(18)30386-3**
- 13 Tunçalp Ö, Were W, MacLennan C, Oladapo O, Gülmezoglu A, Bahl R, et al. Quality of care for pregnant women and newborns – the WHO vision. *BJOG*. 2015;122:1045–9. **Medline:25929823 doi:10.1111/1471-0528.13451**
- 14 World Health Organization. Standards for improving quality of maternal and newborn care in health facilities. Geneva, Switzerland: World Health Organization; 2016. Available: https://cdn.who.int/media/docs/default-source/mca-documents/qoc/quality-of-care/standards-for-improving-quality-of-maternal-and-newborn-care-in-health-facilities.pdf?sfvrsn=3b364d8_4. Accessed: 20 January 2026.
- 15 World Health Organization, United Nations Children's Fund, United Nations Population Fund. Every Woman Every Newborn Everywhere. 2024. Available: <https://ewene.org/>. Accessed: 31 December 2024.
- 16 World Health Organization. Networks of care for maternal and newborn health: implementation guidance 2024. Geneva, Switzerland: World Health Organization; 2024. Available: <https://iris.who.int/bitstream/handle/10665/379031/9789240098633-eng.pdf>. Accessed: 30 December 2024.
- 17 Arksey H, O'Malley L. Scoping studies: towards a methodological framework. *Int J Soc Res Methodol*. 2005;8:19–32. **doi:10.1080/1364557032000119616**
- 18 Peters MDJ, Godfrey CM, Khalil H, McInerney P, Parker D, Soares CB. Guidance for conducting systematic scoping reviews. *Int J Evid-Based Healthc*. 2015;13:141–6. **Medline:26134548 doi:10.1097/XEB.0000000000000050**
- 19 Levac D, Colquhoun H, O'Brien KK. Scoping studies: advancing the methodology. *Implement Sci*. 2010;5:69. **Medline:20854677 doi:10.1186/1748-5908-5-69**
- 20 Tricco AC, Lillie E, Zarin W, O'Brien KK, Colquhoun H, Levac D, et al. PRISMA Extension for Scoping Reviews (PRISMA-ScR): Checklist and Explanation. *Ann Intern Med*. 2018;169:467–73. **Medline:30178033 doi:10.7326/M18-0850**
- 21 Page MJ, McKenzie JE, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD, et al. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *BMJ*. 2021;372:n71. **Medline:33782057 doi:10.1136/bmj.n71**
- 22 Carmone AE, Kalaris K, Leydon N, Sirivansanti N, Smith JM, Storey A, et al. Developing a Common Understanding of Networks of Care through a Scoping Study. *Health Syst Reform*. 2020;6:e1810921. **Medline:33021881 doi:10.1080/23288604.2020.1810921**
- 23 Healthy Newborn Network. Data [dataset]. 2022. Available: <https://www.healthynewbornnetwork.org/numbers/>. Accessed: 20 January 2026.
- 24 Barnett-Page E, Thomas J. Methods for the synthesis of qualitative research: a critical review. *BMC Med Res Methodol*. 2009;9:59. **Medline:19671152 doi:10.1186/1471-2288-9-59**
- 25 Donabedian A. The quality of care. How can it be assessed? *JAMA*. 1988;260:1743–8. **Medline:3045356 doi:10.1001/jama.1988.03410120089033**
- 26 Amoro VA, Abiiro GA, Alatinga KA. Bypassing primary healthcare facilities for maternal healthcare in North West Ghana: socio-economic correlates and financial implications. *BMC Health Serv Res*. 2021;21:545. **Medline:34078379 doi:10.1186/s12913-021-06573-3**
- 27 Gething PW, Johnson FA, Frempong-Ainguah F, Nyarko P, Baschieri A, Aboagye P, et al. Geographical access to care at birth in Ghana: a barrier to safe motherhood. *BMC Public Health*. 2012;12:991. **Medline:23158554 doi:10.1186/1471-2458-12-991**
- 28 Tette EMA, Nartey ET, Nuertey BD, Azusong EA, Akaateba D, Yirifere J, et al. The pattern of neonatal admissions and mortality at a regional and district hospital in the Upper West Region of Ghana; a cross sectional study. *PLoS One*. 2020;15:e0232406. **Medline:32365073 doi:10.1371/journal.pone.0232406**

- 29 Dillip A, Kimatta S, Embrey M, Chalker JC, Valimba R, Malliwah M, et al. Can formalizing links among community health workers, accredited drug dispensing outlet dispensers, and health facility staff increase their collaboration to improve prompt access to maternal and child care? A qualitative study in Tanzania. *BMC Health Serv Res.* 2017;17:416. **Medline:28629475** doi:10.1186/s12913-017-2382-1
- 30 Kroelinger CD, Rice ME, Okoroh EM, DeSisto CL, Barfield WD. Seven years later: state neonatal risk-appropriate care policy consistency with the 2012 American Academy of Pediatrics Policy. *J Perinatol.* 2022;42:595–602. **Medline:34253843** doi:10.1038/s41372-021-01146-y
- 31 Hankins GDV, Clark SL, Pacheco LD, O’Keeffe D, D’Alton M, Saade GR. Maternal mortality, near misses, and severe morbidity: lowering rates through designated levels of maternity care. *Obstet Gynecol.* 2012;120:929–34. **Medline:22996111** doi:10.1097/AOG.0b013e31826af878
- 32 Bolbocean C, Wintermark P, Shevell MI, Oskoui M. Perinatal Regionalization and Implications for Long-Term Health Outcomes in Cerebral Palsy. *Can J Neurol Sci.* 2016;43:248–53. **Medline:26790470** doi:10.1017/cjn.2015.322
- 33 Catalano A, Bennett A, Busacker A, Carr A, Goodman D, Kroelinger C, et al. Implementing CDC’s Level of Care Assessment Tool (LOCATe): A National Collaboration to Improve Maternal and Child Health. *J Womens Health (Larchmt).* 2017;26:1265–9. **Medline:29240547** doi:10.1089/jwh.2017.6771
- 34 Menard MK, Kilpatrick S, Saade G, Hollier LM, Joseph GFJ, Barfield W, et al. Levels of maternal care. *Am J Obstet Gynecol.* 2015;212:259–71. **Medline:25620372** doi:10.1016/j.ajog.2014.12.030
- 35 Chilvers R. Planning Framework for Human Resources for Health for Maternal and Newborn Care [dissertation]. London, UK: London School of Hygiene and Tropical Medicine; 2014. Available: <https://researchonline.lshtm.ac.uk/id/eprint/2124342/>. Accessed: 20 January 2026.
- 36 Cattaneo A, Amani A, Charpak N, De Leon-Mendoza S, Moxon S, Nimbalkar S, et al. Report on an international workshop on kangaroo mother care: lessons learned and a vision for the future. *BMC Pregnancy Childbirth.* 2018;18:170. **Medline:29769056** doi:10.1186/s12884-018-1819-9
- 37 Goldenberg RL, Jones B, Griffin JB, Rouse DJ, Kamath-Rayne BD, Trivedi N, et al. Reducing maternal mortality from preeclampsia and eclampsia in low-resource countries—what should work? *Acta Obstet Gynecol Scand.* 2015;94:148–55. **Medline:25353716** doi:10.1111/aogs.12533
- 38 Simioni AT, Llanos O, Romero M, Ramos S, Brizuela V, Abalos E. [Regionalization of perinatal health care in the province of Santa Fe, Argentina]. *Rev Panam Salud Publica.* 2017;41:e38. Spanish. **Medline:28591333**
- 39 Keene CM, Jalemba A, Murphy GAV, Abuya N, Gathara D, English M, et al. Developing recommendations for neonatal inpatient care service categories: reflections from the research, policy and practice interface in Kenya. *BMJ Glob Health.* 2019;4:e001195. **Medline:30997163** doi:10.1136/bmjgh-2018-001195
- 40 Fawcus S. Practical approaches to managing postpartum haemorrhage with limited resources. *Best Pract Res Clin Obstet Gynaecol.* 2019;61:143–55. **Medline:31103529** doi:10.1016/j.bpobgyn.2019.03.009
- 41 Long Q, Madede T, Parkkali S, Chavane L, Sundby J, Hemminki E. Maternity care system in Maputo, Mozambique: Plans and practice? *Cogent Med.* 2017;4:1412138. doi:10.1080/2331205X.2017.1412138
- 42 Moxon SG, Blencowe H, Bailey P, Bradley J, Day LT, Ram PK, et al. Categorising interventions to levels of inpatient care for small and sick newborns: Findings from a global survey. *PLoS One.* 2019;14:e0218748. **Medline:31295262** doi:10.1371/journal.pone.0218748
- 43 Sharma G, Mathai M, Dickson KE, Weeks A, Hofmeyr GJ, Lavender T, et al. Quality care during labour and birth: a multi-country analysis of health system bottlenecks and potential solutions. *BMC Pregnancy Childbirth.* 2015;15:S2. **Medline:26390886** doi:10.1186/1471-2393-15-S2-S2
- 44 Tappis H, Koblinsky M, Winch PJ, Turkmani S, Bartlett L. Context matters: Successes and challenges of intrapartum care scale-up in four districts of Afghanistan. *Glob Public Health.* 2016;11:387–406. **Medline:26645366** doi:10.1080/17441692.2015.1114657
- 45 Azzuqa A, Makkar A, Machut K. Use of Telemedicine for subspecialty support in the NICU setting. *Semin Perinatol.* 2021;45:151425. **Medline:33992444** doi:10.1016/j.sempperi.2021.151425
- 46 Hall-Barrow J, Hall RW, Burke BLJ. Telemedicine and neonatal regionalization of care - ensuring that the right baby gets to the right nursery. *Pediatr Ann.* 2009;38:557–61. **Medline:19968193** doi:10.3928/00904481-20090918-02
- 47 Handley SC, Srinivas SK, Lorch SA. Regionalization of Care and the Maternal-Infant Dyad Disconnect. *JAMA.* 2019;322:503–4. **Medline:31282951** doi:10.1001/jama.2019.6403
- 48 Harrison WN, Wasserman JR, Goodman DC. Regional Variation in Neonatal Intensive Care Admissions and the Relationship to Bed Supply. *J Pediatr.* 2018;192:73–79.e4. **Medline:28969888** doi:10.1016/j.jpeds.2017.08.028

- 49 Kunz SN, Helkey D, Zitnik M, Phibbs CS, Rigdon J, Zupancic JAF, et al. Quantifying the variation in neonatal transport referral patterns using network analysis. *J Perinatol*. 2021;41:2795–803. **Medline:34035453** doi:10.1038/s41372-021-01091-w
- 50 Narli N, Kırımı E, Uslu S. Turkish Neonatal Society guideline on the safe transport of newborn. *Turk Pediatri Ars*. 2018;53:S18–31. **Medline:31236016** doi:10.5152/TurkPediatriArs.2018.01804
- 51 Soliman Y, Paul R, Pearson K, Alshaikh B, Thomas S, Yusuf K. Neonatal transport services, a cross-sectional study. *J Matern Fetal Neonatal Med*. 2021;34:774–9. **Medline:31084226** doi:10.1080/14767058.2019.1618823
- 52 Bhatta S, Rajbhandari S, Kalaris K, Carmone AE. The Logarithmic Spiral of Networks of Care for Expectant Families in Rural Nepal: A Descriptive Case Study. *Health Syst Reform*. 2020;6:e1824520. **Medline:33253035** doi:10.1080/23288604.2020.1824520
- 53 James A, Endacott R, Stenhouse E. Identifying women requiring maternity high dependency care. *Midwifery*. 2011;27:60–6. **Medline:19837493** doi:10.1016/j.midw.2009.09.001
- 54 Ameyaw EK, Njue C, Tran NT, Dawson A. Quality and women's satisfaction with maternal referral practices in sub-Saharan African low and lower-middle income countries: a systematic review. *BMC Pregnancy Childbirth*. 2020;20:682. **Medline:33176732** doi:10.1186/s12884-020-03339-3
- 55 Marcus JK, Clow SE. Response times of ambulances to calls from midwife obstetric units of the Peninsula Maternal and Neonatal Service (PMNS) in Cape Town. *Curationis*. 2009;32:59–66. **Medline:20225754** doi:10.4102/curationis.v32i1.886
- 56 Busumani W, Mundagowa PT. Outcomes of pregnancy-related referrals from rural health facilities to two central hospitals in Harare, Zimbabwe: a prospective descriptive study. *BMC Health Serv Res*. 2021;21:276. **Medline:33766018** doi:10.1186/s12913-021-06289-4
- 57 Rahman A, Moran A, Pervin J, Rahman A, Rahman M, Yeasmin S, et al. Effectiveness of an integrated approach to reduce perinatal mortality: recent experiences from Matlab, Bangladesh. *BMC Public Health*. 2011;11:914. **Medline:22151276** doi:10.1186/1471-2458-11-914
- 58 Okoroh EM, Kroelinger CD, Lasswell SM, Goodman DA, Williams AM, Barfield WD. United States and territory policies supporting maternal and neonatal transfer: Review of transport and reimbursement. *J Perinatol*. 2016;36:30–4. **Medline:26334399** doi:10.1038/jp.2015.109
- 59 Dosani A, Murthy P, Kassam S, Rai B, Lodha AK. Parental perception of neonatal transfers from level 3 to level 2 neonatal intensive care units in Calgary, Alberta: qualitative findings. *BMC Health Serv Res*. 2021;21:981. **Medline:34535124** doi:10.1186/s12913-021-06967-3
- 60 Daniels AA, Abuosi A. Improving emergency obstetric referral systems in low and middle income countries: a qualitative study in a tertiary health facility in Ghana. *BMC Health Serv Res*. 2020;20:32. **Medline:31924203** doi:10.1186/s12913-020-4886-3
- 61 M'Rithaa DKM, Fawcus SR, De la Harpe M, Korpela M. Development spots in communication during the management of the intrapartum period: An interpretive multiple case study in a developing context. *Afr J Prim Health Care Fam Med*. 2017;9:e1–6. **Medline:28828876** doi:10.4102/phcfm.v9i1.1239
- 62 Rigoli F, Mascarenhas S, Alves D, Canelas T, Duarte G. Tracking pregnant women displacements in Sao Paulo, Brazil: a complex systems approach to regionalization through the emergence of patterns. *BMC Med*. 2019;17:184. **Medline:31570106** doi:10.1186/s12916-019-1416-4
- 63 Fasawe O, Adekeye O, Carmone AE, Dahunsi O, Kalaris K, Storey A, et al. Applying a Client-centered Approach to Maternal and Neonatal Networks of Care: Case Studies from Urban and Rural Nigeria. *Health Syst Reform*. 2020;6:e1841450. **Medline:33270477** doi:10.1080/23288604.2020.1841450
- 64 Staebler S. Regionalized systems of perinatal care: health policy considerations. *Adv Neonatal Care*. 2011;11:37–42. **Medline:21285655** doi:10.1097/ANC.0b013e318206fd5a
- 65 Kornelsen J, Moola S, Grzybowski S. Does Distance Matter? Increased Induction Rates for Rural Women Who Have to Travel for Intrapartum Care. *J Obstet Gynaecol Can*. 2009;31:21–7. **Medline:19208279** doi:10.1016/S1701-2163(16)34049-X
- 66 Gómez López ME. Psychological care guide for women with obstetric complications. *Perinatol Reprod Hum*. 2018;32:85–92.
- 67 Teela KC, Mullany LC, Lee CI, Poh E, Paw P, Masenior N, et al. Community-based delivery of maternal care in conflict-affected areas of eastern Burma: perspectives from lay maternal health workers. *Soc Sci Med*. 2009;68:1332–40. **Medline:19232808** doi:10.1016/j.socscimed.2009.01.033
- 68 Cavallaro FL, Benova L, Wong K, Sheppard P, Faye A, Radovich E, et al. What the percentage of births in facilities does not measure: readiness for emergency obstetric care and referral in Senegal. *BMJ Glob Health*. 2020;5:e001915. **Medline:32201621** doi:10.1136/bmjgh-2019-001915

- 69 Mukuru M, Kiwanuka SN, Gibson L, Ssengooba F. Challenges in implementing emergency obstetric care (EmOC) policies: perspectives and behaviours of frontline health workers in Uganda. *Health Policy Plan.* 2021;36:260–72. **Medline:33515014 doi:10.1093/heapol/czab001**
- 70 O'Malley EG, Popivanov P, Fergus A, Tan T, Byrne B. Maternal near miss: what lies beneath? *Eur J Obstet Gynecol Reprod Biol.* 2016;199:116–20. **Medline:26921477 doi:10.1016/j.ejogrb.2016.01.031**
- 71 Lorch SA, Baiocchi M, Ahlberg CE, Small DS. The differential impact of delivery hospital on the outcomes of premature infants. *Pediatrics.* 2012;130:270–8. **Medline:22778301 doi:10.1542/peds.2011-2820**
- 72 Wright JD, Herzog TJ, Shah M, Bonanno C, Lewin SN, Cleary K, et al. Regionalization of care for obstetric hemorrhage and its effect on maternal mortality. *Obstet Gynecol.* 2010;115:1194–200. **Medline:20502290 doi:10.1097/AOG.0b013e3181df94e8**
- 73 Neogi SB, Malhotra S, Zodpey S, Mohan P. Does facility based newborn care improve neonatal outcomes? A review of evidence. *Indian Pediatr.* 2012;49:651–8. **Medline:22962238 doi:10.1007/s13312-012-0135-0**
- 74 Kern-Goldberger AR, Friedman A, Moroz L, Gyamfi-Bannerman C. Racial Disparities in Maternal Critical Care: Are There Racial Differences in Level of Care? *J Racial Ethn Health Disparities.* 2022;9:679–83. **Medline:33686625 doi:10.1007/s40615-021-01000-z**
- 75 Korst LM, Feldman DS, Bollman DL, Fridman M, El Haj Ibrahim S, Fink A, et al. Cross-sectional survey of California childbirth hospitals: implications for defining maternal levels of risk-appropriate care. *Am J Obstet Gynecol.* 2015;213:527.e1–e12. **Medline:26196455 doi:10.1016/j.ajog.2015.07.014**
- 76 Scott J. Obstetric Transport. *Obstet Gynecol Clin North Am.* 2016;43:821–40. **Medline:27816163 doi:10.1016/j.ogc.2016.07.013**
- 77 Handley SC, Passarella M, Srinivas SK, Lorch SA. Identifying individual hospital levels of maternal care using administrative data. *BMC Health Serv Res.* 2021;21:538. **Medline:34074286 doi:10.1186/s12913-021-06516-y**
- 78 Isayama T. The clinical management and outcomes of extremely preterm infants in Japan: past, present, and future. *Transl Pediatr.* 2019;8:199–211. **Medline:31413954 doi:10.21037/tp.2019.07.10**
- 79 Blondel B, Papiernik E, Delmas D, Künzel W, Weber T, Maier RF, et al. Organisation of obstetric services for very preterm births in Europe: Results from the MOSAIC project. *BJOG.* 2009;116:1364–72. **Medline:19538415 doi:10.1111/j.1471-0528.2009.02239.x**
- 80 Kunz SN, Phibbs CS, Profit J. The changing landscape of perinatal regionalization. *Semin Perinatol.* 2020;44:151241. **Medline:32248957 doi:10.1016/j.semperi.2020.151241**
- 81 Brantley MD, Davis NL, Goodman DA, Callaghan WM, Barfield WD. Perinatal regionalization: a geospatial view of perinatal critical care, United States, 2010–2013. *Am J Obstet Gynecol.* 2017;216:185.e1–10. **Medline:27773712 doi:10.1016/j.ajog.2016.10.011**
- 82 Clapp MA, Srinivas SK, Kozhimannil KB, Grobman WA, Kaimal AJ. Comparison of Maternal and Neonatal Subspecialty Care Provision by Hospital. *Am J Perinatol.* 2023;40:333–40. **Medline:33878766 doi:10.1055/s-0041-1727223**
- 83 Barnea ER, Nicholson W, Theron G, Ramasauskaite D, Stark M, Albini SM, et al. From fragmented levels of care to integrated health care: Framework toward improved maternal and newborn health. *Int J Gynaecol Obstet.* 2021;152:155–64. **Medline:33341938 doi:10.1002/ijgo.13551**
- 84 Zahn CM, Remick A, Catalano A, Goodman D, Kilpatrick SJ, Menard MK. Levels of Maternal Care Verification Pilot: Translating Guidance Into Practice. *Obstet Gynecol.* 2018;132:1401–6. **Medline:30399104 doi:10.1097/AOG.0000000000002952**
- 85 Suehara N. Cross-regional perinatal healthcare systems: Efforts in the Greater Kansai Area, Japan. *JMAJ.* 2010;53:86–90.
- 86 Kc A, Bhandari A, Pradhan YV, Kc NP, Upreti SR, Thapa K, et al. State of maternal, newborn and child health programmes in Nepal: what may a continuum of care model mean for more effective and efficient service delivery? *J Nepal Health Res Counc.* 2011;9:92–100. **Medline:22929837**
- 87 Boerma T, Campbell OMR, Amouzou A, Blumenberg C, Blencowe H, Moran A, et al. Maternal mortality, stillbirths, and neonatal mortality: a transition model based on analyses of 151 countries. *Lancet Glob Health.* 2023;11:e1024–31. **Medline:37349032 doi:10.1016/S2214-109X(23)00195-X**