G Model VIRUS 95485 1-8

Virus Research xxx (2011) xxx-xxx



Contents lists available at ScienceDirect

Virus Research



journal homepage: www.elsevier.com/locate/virusres

Please cite this article in press as: Quan, P.-L., et al., Genetic characterization of K13965, a strain of Oak Vale virus from Western Australia. Virus

Genetic characterization of K13965, a strain of Oak Vale virus from Western Australia

Phenix-Lan Quan^a, David T. Williams^b, Cheryl A. Johansen^c, Komal Jain^a, Alexandra Petrosov^a, Sinead M. Diviney^b, Alla Tashmukhamedova^a, Stephen K. Hutchison^d, Robert B. Tesh^e, John S. Mackenzie^{b,f}, Thomas Briese^{a,g,*}, W. Ian Lipkin^{a,g,h,i}

^a Center for Infection and Immunity, Columbia University, New York, NY, USA

^b School of Biomedical Sciences, Curtin University, Perth, WA, Australia

d 454 Roche Life Sciences, Branford, CT, USA

11 ^f Burnet Institute, Melbourne, Australia

^g Department of Epidemiology, Mailman School of Public Health, Columbia University, New York, NY, USA 12

h Department of Pathology, College of Physicians and Surgeons, Columbia University, New York, NY, USA 13

14 ⁱ Department of Neurology, College of Physicians and Surgeons, Columbia University, New York, NY, USA

ARTICLE INFO

15

16 17

31

33

34

37

38

Article history: 18 Received 12 April 2011 19 Received in revised form 17 June 2011 20 Accepted 21 June 2011 21 Available online xxx 22 23 Keywords: Rhabdovirus 24 Oak Vale virus 25 Anopheles annulipes 26 27 Culex edwardsi Virus phylogenetics 28 29 Australia

Molecular characterization 30

1. Introduction

Viruses with nonsegmented, single-stranded negative sense RNA genomes are subsumed in the order Mononegavirales that currently includes the four families Bornaviridae, Filoviridae, Paramyxoviridae and Rhabdoviridae. The family Rhabdoviridae comprises viruses of fish in the genus Novirhabdovirus and arthropod-borne viruses that infect plants in the genera Nucleorhabdovirus and Cytorhabdovirus. The remaining genera Ephemerovirus, Lyssavirus and Vesiculovirus include both arthropod-borne and non vector-39 borne pathogens of mammals, birds, reptiles and fish that include 40

* Corresponding author at: Center for Infection and Immunity, Mailman School of Public Health, Columbia University, 722 West 168th Street, New York, NY 10032, USA. Tel.: +1 212 342 9031: fax: +1 212 342 9044.

E-mail addresses: thomas.briese@columbia.edu, tb2047@columbia.edu, tbriese@ymail.com (T. Briese).

0168-1702/\$ - see front matter © 2011 Published by Elsevier B.V. doi:10.1016/j.virusres.2011.06.021

Res. (2011), doi:10.1016/j.virusres.2011.06.021

ABSTRACT

K13965, an uncharacterized virus, was isolated in 1993 from Anopheles annulipes mosquitoes collected in the Kimberley region of northern Western Australia. Here, we report its genomic sequence, identify it as a rhabdovirus, and characterize its phylogenetic relationships. The genome comprises a P' (C) and SH protein similar to the recently characterized Tupaia and Durham viruses, and shows overlap between G and L genes. Comparison of K13965 genome sequence to other rhabdoviruses identified K13965 as a strain of the unclassified Australian Oak Vale rhabdovirus, whose complete genome sequence we also determined. Phylogenetic analysis of N and L sequences indicated genetic relationship to a recently proposed Sandjima virus clade, although the Oak Vale virus sequences form a branch separate from the African members of that group.

© 2011 Published by Elsevier B.V.

01

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57

significant threats to agriculture and public health, such as vesicular stomatitis Indiana virus (VSIV) and rabies virus (RABV) (Kuzmin et al., 2009). In addition, more than 100 viruses are tentatively assigned to this family based on their characteristic bullet-shaped virion morphology, or via serologic cross-reactivity with one or more of the classified members of the family (Tordo et al., 2005).

Rhabdoviruses have genomes that range in approximate size from 11,000 to 16,000 nucleotides (nt). The genome organization of many classic rhabdoviruses, as represented by RABV, has been considered to be one of the simplest in the viral kingdom with only 5 genes: the nucleoprotein (N), phosphoprotein (P), matrix protein (M), glycoprotein (G) and polymerase (L) in the order 3'-N-P-M-G-L-5' (Kuzmin et al., 2009). However, recent molecular analyses of novel rhabdoviruses indicated considerable genetic variability and more complex genomes, with up to 10 additional open reading frames (ORF) located within or interposed between the five core genes, many still without known function. In addition to overlapping ORFs characterized within the P gene of members

^c Discipline of Microbiology and Immunology, University of Western Australia, Nedlands, WA, Australia

^e Department of Pathology, University of Texas Medical Branch, Galveston, TX, USA 10

G Model VIRUS 95485 1-8

2

59

81

84

97

90

100

101

102

103

104

105

106

107

108

109

110

111

112

113

114

115

116

117

118

119

120

P.-L. Quan et al. / Virus Research xxx (2011) xxx-xx

of the Vesiculovirus genus (C and C'; Kretzschmar et al., 1996; Peluso et al., 1996; Spiropoulou and Nichol, 1993), and in unas-60 signed rhabdoviruses from vertebrates and dipterans (Allison et al., 61 2010; Gubala et al., 2010; Nishizawa et al., 1997; Quan et al., 62 2010; Schutze et al., 1999; Springfeld et al., 2005; Tao et al., 2008; 63 Zhu et al., 2011), ORFs that overlap within N, M or G genes have 64 also been identified (Gubala et al., 2008; Jayakar and Whitt, 2002; 65 Tao et al., 2008). One to four ORFs located between the P and 66 M genes are reported for plant-infecting rhabdoviruses (Dietzgen 67 et al., 2006; Huang et al., 2003; Reed et al., 2005; Revill et al., 2005; 68 Scholthof et al., 1994; Tanno et al., 2000), the unassigned Drosophila 69 sigma viruses (Longdon et al., 2010; Teninges et al., 1993), and 70 the Culicoides-transmitted Ngaingan (NGAV) and Wongabel viruses 71 (WOGV) (Gubala et al., 2008, 2010). In addition, an ORF located 72 between the M and G genes has been identified in the unassigned 73 NGAV, Durham (DURV), and Tupaia viruses (TUPV) (Allison et al., 74 2010; Gubala et al., 2010; Springfeld et al., 2005). All members of 75 the genus Novirhadovirus, the nucleorhabdovirus rice yellow stunt 76 virus (RYSV) and the unassigned coastal plains virus (CPV, Genbank 77 accession GQ294473) have a unique ORF located between G and L 78 genes, whereas some members of the genus Ephemerovirus contain 79 80 ORFs potentially resulting from gene duplication at that junction (Huang et al., 2003; McWilliam et al., 1997; Morzunov et al., 1995; Wang et al., 1994). Additional genes located between the N and P 82 genes have not yet been reported. 83

Arboviruses of medical importance have been monitored in northern Western Australia (WA) since the 1970s through serol-85 ogy and annual surveys of mosquitoes during the late wet season in 86 major towns and communities across the Kimberley region (Broom 87 et al., 2003; Liehne et al., 1981). During the course of those surveil-88 lance campaigns, a novel virus, designated K13965, was isolated. 89 After K13965 failed to react with a panel of monoclonal antibodies 90 raised to alphaviruses and flaviviruses common in WA(Broom et al., 91 1998), and did not yield amplification products using primers for 92 the detection of common arboviruses, unbiased high-throughput 93 pyrosequencing (UHTS) was employed for genetic characterization. 94 Here we report the complete genome sequence of K13965, and its 95 identification as a rhabdovirus that represents a strain of Oak Vale 96 virus (OVRV), for which we also report full genomic sequence.

2. Materials and methods

2.1. Virus collection and culture

K13965 was isolated from a pool of female Anopheles annulipes s.l. mosquitoes collected in May 1993 from a trapping site located approximately 3 km north of Kununurra in the Kimberley region of WA (Fig. 1). Mosquitoes were collected in an annual survey of mosquitoes and arboviruses across northern WA by the University of Western Australia Arbovirus Surveillance and Research Laboratory (Broom et al., 2002; Johansen et al., 2009a). Mosquitoes were processed for virus isolation by inoculation of C6/36 cells, followed by serial passage of cell culture supernatant in PSEK and Vero cells (Johansen et al., 2000; Lindsay et al., 1993). Infected cultures were monitored by microscopic examination for cytopathic effect. Culture supernatants were harvested, serially passaged, and used for subsequent studies. Virus identification using infected cell monolayers was attempted by enzyme immunoassay using a panel of monoclonal antibodies to Australian arboviruses (Broom et al., 1998), and by reverse transcription-polymerase chain reaction (RT-PCR) assays on nucleic acid extracts using a panel of primer sets targeting alphaviruses, flaviviruses, orbiviruses and bunyaviruses (primer sequences available upon request).

OVRV strain CSIRO-1342 was obtained from the World Reference Center for Emerging Viruses and Arboviruses collection at the



Fig. 1. Map of Australia by states and territories showing the geographic locations where OVRV K13965 and CSIRO-1342 were isolated. K13965 was isolated from Anopheles annulipes s.l. collected 3 km north of Kununurra. OVRV CSIRO-1342 was isolated from Culex edwardsi collected at Peachester. Another isolation of OVRV has been reported from Aedes vigilax collected at Darwin, ACT, Australian Capital Territory; NSW, New South Wales; NT, Northern Territory; QLD, Queensland; SA, South Australia; TAS, Tasmania; VIC, Victoria; WA, Western Australia.

University of Texas Medical Branch. The isolation of OVRV strain CSIRO-1342 was reported from Peachester, Queensland, in 1981/82 from truck-trapped Culex sp. mosquitoes, and in 1984 from Culex edwardsi mosquito pools collected during an ephemeral fever outbreak (Calisher et al., 1989; Cybinski and Muller, 1990; Muller and Standfast, 1986) (Fig. 1).

2.2. Unbiased high-throughput sequencing (UHTS)

Culture supernatant from infected PSEK cells was clarified from cell debris by centrifugation and total RNA was extracted from the clarified supernatant for UHTS (Trizol-LS, Invitrogen, Carlsbad, CA, USA). RNA (0.5 µg) was DNase I-digested (DNA-free; Ambion, Austin, TX, USA) and reverse transcribed using Superscript II (Invitrogen) with random octamer primers linked to an arbitrary, defined 17-mer primer sequence. The cDNA was RNase H treated prior to random amplification by PCR using AmpliTaq (Applied Biosystems, Foster City, CA, USA) and a primer mix including the octamerlinked 17-mer sequence primer and the defined 17-mer sequence primer in a 1:9 ratio (Quan et al., 2007). Products >70 bp were purified (MinElute, Qiagen, Hilden, Germany) and ligated to linkers for sequencing on a GSL FLX Sequencer (454 Life Sciences, Branford, CT, USA) (Margulies et al., 2005). After trimming primer sequences and eliminating highly repetitive elements, sequences were clustered and assembled into contiguous fragments (contigs) for comparison by the basic local alignment search tool (blast; Altschul et al., 1990) to the Genbank database at nucleotide (nt) and deduced amino acid (aa) level, applying blastn and fastx algorithms.

2.3. Specific RT-PCR and quantitative real time RT-PCR

Multiple primer sets were designed based on sequences obtained through UHTS. The draft genome sequence was validated by sequencing overlapping PCR products that covered the entire genome. Genomic termini were characterized by RACE (Invitrogen). RNA was transcribed with Superscript II (Invitrogen) using random hexamer priming. PCR primers were applied at 0.2 µM concentration with 1 µL cDNA and Platinum Tag DNA polymerase (Invitrogen). Products were purified (QIAquick PCR purification kit; Qiagen) and directly dideoxy-sequenced on both strands (Genewiz, South Plainfield, NJ, USA).

Primers and probe for quantitative real-rime PCR were selected within the L gene using Geneious software (Biomatters, Auckland, New Zealand): K13965-fwd 5'-TGGAGGAGACCATGACCAGCACA, 147

148

149

150

151

152

153

154

155

156

157

158

159

160

121

122

ARTICLE IN PRESS



С

Leader **UGC**UCUUUU**U**GUUUUACAAAAGUACAGCUGGUUUGUUAGCUGUGUGCAGUUGGUUC<u>UAC</u> |||| ||||||| Trailer **ACG**AAAAAA**A**CCAAUCCAGAUGUUCUCCCCCCAUCACCGGAUUCACCAACGUUUUUUUCAUGUG<u>UUA</u>

Fig. 2. K13965 genome organization. (A) Schematic organization of the K13965 genome. The bar represents the 11,120 nucleotide (nt) antigenome. The eight open arrows indicate the position of the open reading frames (ORFs) for putative N, P, P', P'', M, SH, G, and L proteins. (B) Transcription initiation, intergenic and transcription termination/polyadenylation sequences. The start and stop codon for each gene is in bold, the intergenic region is underlined. (C) Complementarity between the 3' leader and the 5' trailer sequences. Bases conserved among rhabdoviruses known to infect vertebrates are shown in bold. Start and stop codons are underlined.

K13965-rev 5'-GCTCAGACAGTTGGCTATGTT AGGAAG, K13965-161 probe 5'-FAM-AGCAAGAGGTATGCAAGAATGTGCA-TAMRA. A cal-162 ibration standard was generated by cloning a 102 nt genomic 163 fragment (pGEM-T Easy; Promega, Madison, WI, USA). Assays were 164 performed in duplicate using a StepOnePlus Real-time PCR mix 165 (Applied Biosystems), and a standard cycling profile with 45 cycles 166 in a volume of 25 µL, containing random hexamer-primed cDNA, 167 300 nM primer (each) and 200 nM probe. 168

169 2.4. Phylogenetic analysis

Phylogenetic trees were constructed based on aa sequences of L 170 (970 aa, n = 49, or conserved block III, 156 aa, n = 77) and N (325 aa, n = 77)171 n = 60). Sequences were aligned by the MUSCLE 3.6 software algo-172 rithm (Edgar, 2004) and manually adjusted by using MEGA 4.0 173 174 software (Tamura et al., 2007) after terminal regions with low align-175 ment confidence were clipped. Phylogenetic trees were generated by applying the Bayesian method of the MrBayes software package 176 (v 3.1.2) (Huelsenbeck and Ronquist, 2001), using a Whelan and 177 Goldman (WAG) evolutionary model of aa replacement (Whelan 178 and Goldman, 2001) with a gamma distribution of rate variation 179 among sites. Chains were run for 10 million generations and sam-180 pled every 100th generation, applying a 25% burn-in rate at which 181 point all parameters had converged. 182

183 2.5. Protein analysis

Similarity to protein families and prediction of functional 184 protein domains were obtained through sequence anal-185 yses with PFAM (http://pfam.sanger.ac.uk/) and PROSITE 186 (http://ca.expasy.org/prosite). Predictions of physico-chemical 187 properties of deduced proteins of K13965 (molecular 188 weight (MW), isoelectric point (pI), grand average of 189 hydropathicity (GRAVY)) were generated 190 by using the

Res. (2011), doi:10.1016/j.virusres.2011.06.021

Protparam tool (http://expasy.ch/tools/). Phobius software (http://www.ebi.ac.uk/Tools/phobius/) was used to predict signal peptides and protein topology. Predictions of N-glycosylation and phosphorylation sites were derived with the respective algorithms of the Center for Biological Sequence Analysis (http://www.cbs.dtu.dk/services/). Amino acid sequence identity and similarity were calculated by applying the Needleman algorithm with an EBLOSUM30 substitution matrix (gap open/extension penalties of 10/2 for aa alignments; EMBOSS (Rice et al., 2000)) and a Perl script to compile the results for all comparisons.

3. Results

Please cite this article in press as: Quan, P.-L., et al., Genetic characterization of K13965, a strain of Oak Vale virus from Western Australia. Virus

3.1. K13965 genome organization

The complete K13965 genomic sequence was determined using infected cell culture supernatant. UHTS yielded approximately 112,800 sequence reads with a mean length of 245 nt. After primer trimming and assembly, 6 contigs ranging in size from 96 to 5362 nt (14,363 sequence reads, mean length 353 nt) showed homology to rhabdovirus sequences in the NCBI non-redundant database (http://www.ncbi.nlm.nih.gov/Genbank). The assembled sequence covered full-length genome sequence, except for leader and trailer regions. Genomic termini were subsequently characterized by RACE. The generated draft sequence was confirmed by overlapping PCR across the whole genome (Genbank accession JF705877).

The negative-sense genome sequence of K13965 comprises 11,220 nt; the antigenome encodes at least eight ORFs (Fig. 2A). In addition to the five core rhabdoviral genes, K13965 includes an additional ORF (SH) located between the M and G genes, and two putative ORFs (P', P'') in overlapping reading frames within the P gene to result in a genome organization 3'-Leader-N-P-(P', P'')-M-SH-G-L-Trailer-5'. Putative transcription initiation (3'-CUGU) and

191

192

193

3

212

213

214

215

216

217

218

219

220

221

222

223

224

225

226

227

228

220

230

231

232

233

234

235

236

237

28

P.-L. Quan et al. / Virus Research xxx (2011) xxx-xx

termination (3'-GUACU₇) signals are recognized flanking the identified ORFs, resulting in six transcription units (Fig. 2A and B). Gene junctions show the usual separation of the upstream termination signal by 1-2 nt from the downstream initiation signal, except at the G-L junction. The intergenic regions consist of a dinucleotide (GA) at the N-P and SH-G junctions, whereas a single nucleotide (G or A) is present at the P-M and M-SH junctions (Fig. 2B). In contrast, the termination/polyadenylation signal of G is located 2 nt downstream of the Linitiation sequence, resulting in overlap of the G and L genes without overlap of their ORFs.

A common feature of rhabdovirus genomes is partial complementary of their 3'- and 5'-termini (Whelan et al., 2004). The 3'-leader and the 5'-trailer sequences of K13965 comprise 39 and 51 nt, respectively; with eleven of the fifteen terminal nt being complementary (Fig. 2C).

3.2. Open reading frames

Features of the ORFs identified in the K13965 antigenome are 238 summarized in Table 1. ORF-1 is compatible with nucleoproteins 239 240 (N) of other rhabdoviruses by location, and contains a highly conserved aa motif S289 PYS that is proposed to be involved in RNA-241 binding in vertebrate rhabdoviruses (Crysler et al., 1990; Tordo 242 et al., 1986). Alignment with other N sequences also indicates 243 conserved aa identified as characteristic of rhabdoviruses that 244 infect vertebrates (Kuzmin et al., 2006). ORF-II contains several 245 potential phosphorylation sites, including 2 tyrosine, 24 serine, and 246 7 threonine phosphorylation sites as anticipated for a phospho-247 protein (P), the cofactor of the active viral transcriptase/replicase 248 complex. In addition, two non-overlapping ORFs in the +1 frame 249 of the P gene may encode small acidic proteins designated P' and 250 P'' (Table 1 and Fig. 2). P' is located close to the 5'-end of the P ORF, 251 similar to C proteins identified in VSIV and vesicular stomatitis 252 New Jersey virus (VSNJV) (Peluso et al., 1996; Spiropoulou and 253 Nichol, 1993). Putative start codons for P' and P" are 65 nt and 254 758 nt downstream of the P start codon. Both putative proteins 255 show no significant similarity to known proteins in the PFAM or 256 PROSITE databases. ORF-III commonly codes for a matrix protein 257 (M). The M analog of K13965 includes potential serine (6 sites) 258 259 and threonine (1 site) phosphorylation sites, and no sequence similarity to other known proteins is detected. K13965 ORF-IV 260 includes 2 hydrophobic aa clusters located at the N terminus 261 and the center of the sequence, resulting in a grand average of 262 hydropathicity of +0.595, analogous to the small hydrophobic (SH) 263 proteins recently identified for TUPV and DURV (Allison et al., 264 2010; Springfeld et al., 2005). Phobius analysis indicated a signal 265 peptide (aa 1-23) but no transmembrane anchor, suggesting that 266 a soluble protein may be generated. Despite the limited primary 267 sequence conservation among glycoproteins (G) of rhabdoviruses, 268 269 general structural features including glycosylation sites, and cysteine residues are commonly conserved (Coll, 1995; Walker and 270 Kongsuwan, 1999). Accordingly, ORF-V of K13965 shows features 271 of type I glycoproteins, including an N-terminal signal peptide (aa 272 1-17), an ectodomain (aa 18-501), a transmembrane domain (aa 273 502–521) followed by a short cytoplasmic tail (aa 522–531), two 274 potential N-glycosylation sites (N₃₇₈KTL and N₄₀₅GTT), and four-275 teen cysteine residues including 7 of the 12 cysteines conserved 276 among animal rhabdoviruses (CI₄₀, CIII₈₁, CV₁₁₉, CVI₁₅₈, CVII₁₆₃, 277 CXI₃₀₉, CXII₃₄₂; Walker and Kongsuwan, 1999). The last and 278 largest ORF-VI is predicted to translate into a 2051 aa (231 kDa) 279 L-polymerase. The L-polymerase of K13965 contains the con-280 served residues present in L-polymerase of negative strand RNA viruses (Poch et al., 1990). In addition, all four highly conserved 282 283 motifs A through D of block III (Poch et al., 1989, 1990) are con-284 served in L of K13965 (motif A (A₅₇₇NHMDYSKWNNHQR);

motif B (A₆₄₇CWRGQAGGLEGLRQKGWTITSLLMI); motif C (T₆₈₅LAQGDNQIV); motif D (Y₇₅₈RGNLCNPKSKRY)).

3.3. Sequence comparison of K13965 with OVRV CSIRO-1342 and other rhabdoviruses

In an independent approach we also determined the complete genomic sequence of another unassigned Australian rhabdovirus, OVRV strain CSIRO-1342. As with K13965, culture supernatant of CSIRO-1342 was analyzed by UHTS, sequence gaps filled with specific PCR and RACE, and the draft sequence finally confirmed through re-sequencing by overlapping PCR (Genbank accession JF705876). The total RNA preparations from CSIRO-1342 and K13965 that were used for UHTS were subsequently evaluated by a real-time RT-PCR assay developed based on the consensus genome sequences. The quantitative OVRV assay indicated virus loads of 1.5×10^5 copies/ng total RNA for CSIRO-1342, and 3.5×10^5 copies/ng for K13965 (Table 2). 02

Like K13965, the negative-sense genome of OVRV comprises 11,220 nt with a comparable genome organization. Furthermore, identical putative transcription initiation, termination and intergenic regions were identified for CSIRO-1342, except for a change at the M termination sequence (3'-GCACU₇). As in K13965 the G and L genes overlap. The putative P' ORFs of K13965 and CSIRO-1342 show high nt sequence conservation of 98.6%, and are identical at the aa level. However, the putative P" start and stop codons are positioned differently in the same +1 frame, generating proteins of 67 aa for K13965, or 48 aa for CSIRO-1342. Comparison of OVRV CSIRO-1342 and K13965 genomes indicated 93% nt identity. The P (92%) and SH (93%) ORFs show the least, and N (100%) the highest aa sequence identity (Suppl. Table 1).

Comparison of individual K13965 ORFs to homologous proteins of other representative rhabdoviruses for which complete sequence is available indicated that K13965 shares at most 36% aa identity for L with TUPV and DURV; P and M show less then 17% identity with the analyzed sequences, and identities for N ranged between 15 and 30% (Suppl. Table 2).

3.4. Phylogenetic analyses

Phylogenetic trees constructed from partial L aa sequences of K13965, CSIRO-1342 and representative rhabdoviruses suggest that K13965 and CSIRO-1342 are closely related to, but distinct from, members of the recently proposed Sandjimba group (Fig. 3A) (Dacheux et al., 2010). Analysis of N aa sequences similarly grouped K13965 and CSIRO-1342 with the available Sandjimba group viruses KOLV and SJAV (Fig. 3B), but also showed a relationship to TUPV and DURV. Only the latter relationship could be confirmed by using nearly full-length L sequence due to a lack of sequence information for the other viruses (Fig. 3C).

4. Discussion

K13965, an uncharacterized Australian virus isolate, was genetically identified as a strain of OVRV by comparison of its genome to the full length genomic sequence that we obtained for the Australian OVRV CSIRO-1342, which matched the previously available partial sequence for CSIRO-1342 (Genbank accession GQ294474). The K13965 genome, in addition to 5 core proteins typical for rhabdoviruses, encodes a small hydrophobic (SH) protein, and putative P' (C) and P" proteins that may be expressed from alternative ORFs contained within the P gene. Like sigma viruses in Drosophila spp. (Longdon et al., 2010; Teninges et al., 1993), K13965, isolated from mosquitoes has overlap between the G and L gene without ORF overlap.

286

285

287 288 289

291

202

203

204

295

296

297

298

299

300

301

302

303

304

305

306

307

308

309

310

311

312

313

314

315

316

317

318

319

320

321

322

323

324

325

326

327

328

329

330

331

332

333

334

335

336

337

338

339

340

341

342

343

Please cite this article in press as: Quan, P.-L., et al., Genetic characterization of K13965, a strain of Oak Vale virus from Western Australia. Virus Res. (2011), doi:10.1016/j.virusres.2011.06.021

344

345

346

347

348

349

350

351

352

353

354

355

356

357

358

ARTICLE IN PRESS

P.-L. Quan et al. / Virus Research xxx (2011) xxx-xx

Table I	
Characteristics of ORFs identified in the K13965 ge	nome

ORF	Gene length (nt)	ORF length (nt)	Protein length (aa)	Protein calculated MW (kDa)	pI	GRAVY
N	1390	1350	449	50.8	6.11	-0.302
Р	1198	1170	389	41.4	5.58	-0.569
P'	_	276	91	10.4	6.10	-0.300
P''	_	204	67	7.6	6.83	-0.367
Μ	528	504	167	19.2	9.73	-0.140
SH	212	183	60	6.8	6.50	0.595
G	1637	1596	531	60.7	8.89	-0.296
L	6176	6156	2051	232	8.57	-0.203

pI: isoelectric point; MW: molecular weight; GRAVY: grand average of hydropathicity.

Within the order *Mononegavirales*, SH and C proteins were initially identified in paramyxoviruses (Bellini et al., 1985; Curran and Kolakofsky, 2008; Elango et al., 1989). Paramyxovirus C proteins have been shown to inhibit interferon signaling (Nagai and Kato, 2004), and SH proteins to inhibit TNF alpha production (Li et al., 2011) and apoptosis (Wilson et al., 2006). Protein products coded in analogous genome locations to paramyxovirus C proteins have been identified in the rhabdoviruses VSIV and VSNJV (Kretzschmar et al., 1996; Peluso et al., 1996; Spiropoulou and Nichol, 1993), and products comparable to SH proteins were identified in TUPV (Springfeld et al., 2005) and predicted for DURV (Allison et al., 2010). The functions of these proteins in rhabdoviruses are unknown although it has been hypothesized that C proteins may play a role in viral pathogenesis and enhance transcriptional activity (Peluso et al., 1996).

The putative P' (C) of K13965 is larger than those reported for 359 members of the genus Vesiculovirus, but smaller than P'(C) reported 360 for TUPV or DURV (Suppl. Table 3). The P' (C) of K13965 is pre-361 dicted to have an acidic pI whereas other rhabdoviral P'(C) proteins, 362 with the exception of that of DURV, have a basic pI >9. P'(C) of 363 K13965 shows no recognized sequence homology to other reported 364 P' (C), except for that of OVRV CSIRO-1342. In addition, a small P" 365 ORF located downstream of P' was identified in K13965, resem-366 bling the P gene organization of NGAV, which is reported to also 367 code for two putative non-overlapping small ORFs (Gubala et al., 368 2010). Although a P" ORF is present also in CSIRO-1342, it is not 369 conserved with respect to K13965; different start and stop codons 370 predict a smaller protein for CSIRO-1342. This lack of conservation 371 372 between the two isolates indicates a lower selective pressure on these sequences compared to the other coding sequences. 373

Genome organization of representative rhabdoviruses.

Table 2

Virus	Genus	Genome structure	Host/vector	Length (nt)	Genbank accession no.
K13956	UC	N-P (P', P")- M -SH- G-L	Mosquitoes	11,120	JF705877
RABV	Lyssavirus	N-P-M-G-L	Bat	11,923	EU293116
VSIV	Vesiculovirus	N-P(C)-M-G-L	Bovine	11,155	AF473865
SIGMAV	UA	N-P-X-M-G-L	Drosophila melanogaster	12,625	GQ375258
SCRV	UC	N-P (C)- M (Ms)- G-L	Fish	11,545	NC_008514
IHNV	Novirhabdovirus	N-P-M-G-NV-L	Fish	11,131	NC_001652
TUPV	Tentative Vesiculovirus	N-P(C)-M-SH-G-L	Tree shrew	11,440	NC_007020
RYSV	Nucleorhabdovirus	N-P-3-M-G-6-L	Plant	14,042	NC_003746
MFSV	Nucleorhabdovirus	N-P-3-4-M-G-L	Plant	13,782	NC_005974
TIBV	UA	N-P-M-U1-U2-G(G')-U3-L	Cattle, dipteran	13,298	GQ294472
CPV	UC	N-P (P')- M -U1-U2- G -U3-L	Livestock	13,203	GQ294473
WOGV	UC	N(U4)-P-U1-U2-U3-M-G(U5)-L	Biting midges	13,196	NC_011639
NCMV	Cytorhabdovirus	N-P-3-4-5-6-M-G-L	Plant	13,222	NC_002251
ARV	Ephemerovirus	N-P-M-G -G _{NS} - $(\alpha 1/\alpha 2)$ - β -L	Bovine	NA	U05987
					U10363
					L09206
BEFV	Ephemerovirus	$N-P(P')-M-G-G_{NS}-(\alpha 1/\alpha 2/\alpha 3)-(\beta/\gamma)-L$	Bovine	14,900	NC_002526
NGAV	UA	N-P (P'1 P'2)-(U1/U2)-U3-M-U4-G-Gns-(U5/U6)-U7-L	Biting midges	15 764	NC 013955

UA: unassigned rhabdovirus; UC: unclassified virus (not found in the VIIIth report of the International Committee on taxonomy viruses). The 5 typical core genes N, P, M, G and L present in all rhabdoviruses are indicated in bold. ORFs translated from polycistronic RNA are indicated in parenthesis.

Both OVRV K13965 and CSIRO-1342 encode a putative hydrophobic SH protein of smaller size than those of TUPV and DURV (Suppl. Table 4) that has no sequence homology with these SHs. TUPV and DURV SHs are predicted to adopt a type I, or type II, transmembrane topology, respectively (Allison et al., 2010; Springfeld et al., 2005). In contrast, SH of OVRV is not predicted to be a transmembrane protein. Interestingly, in NGAV an ORF (U4; Gubala et al., 2010) is present in the same genomic position as SH proteins that also likely generates a soluble but not hydrophobic protein (NC_013955; Phobius). No aa conservation was found in the N-terminal half of the ORF or the entire aa sequence. We also found no conservation of the leucine residues with respect to TUPV and DURV in the SH of K13965 or CSIRO-1342. Together with TUPV and DURV (isolated from mammal and bird, respectively), the mosquito-borne OVRV represents the third example of a rhabdovirus encoding a P' (C) and SH protein.

5

374

375

376

377

378

379

380

381

382

383

384

385

386

387

388

389

390

391

392

393

394

395

396

397

398

399

400

401

402

403

The geographic region from where K13965-infected *An. annulipes* s.l. mosquitoes were collected in 1993 is an enzootic focus for arboviruses, as the presence of the Ord River irrigation area enables mosquito species to breed continuously (Broom et al., 2002, 2003; Liehne et al., 1976; Mackenzie and Broom, 1999). The high level of rainfall during the 1992/1993 wet season was associated with an increase in arbovirus activity in that area (Broom et al., 1997). An isolate of the recently identified Stretch Lagoon orbivirus was also recovered from mosquitoes collected from the Ord River area in 1993 (D.T.W. and C.A.J., unpublished), suggesting that environmental conditions were suitable for heightened arbovirus activity at that time. OVRV CSIRO-1342 was isolated in the early 1980s from *Cx. edwardsi* and *Culex* spp. mosquitoes from southeast Queensland (Calisher et al., 1989; Cybinski and Muller,

Please cite this article in press as: Quan, PL., et al., Genetic cl	characterization of K13965, a strain of Oak Vale virus from Western Austr	alia. Viru
Res. (2011), doi:10.1016/j.virusres.2011.06.021		

6

ARTICLE IN PRESS

P.-L. Quan et al. / Virus Research xxx (2011) xxx-xxx



Fig. 3. Phylogenetic relationships of K13965 and CSIRO-1342 with representative rhabdoviruses derived from partial L(156 aa; A), N(325 aa; B) and full Lamino acid sequence (970 aa; C). The geographic origin (*Africa, #Australia) is indicated for K13965, CSIRO-1342 and members of the Sandjimba and Almpiwar groups, K13965 is indicated by a circled arrowhead. Analysis was performed using a Bayesian method applying a WAG model of amino acid replacement with a gamma distribution of rate variation among sites for 10 million generations (with a 25% burn-in). Support for each node is provided by BPP (Bayesian posterior probability) (>0.8). Branch lengths are drawn to scale and the trees were mid-point rooted. Amino acid sequences used were: ABLV, Australian bat lyssavirus (L:NP.478343; N:AAD01267); ALMV, Almpiwar virus (L:AY854645); ARAV, Aravan virus (L:ABV03822; N:Q6X1D8); ARV, Adelaide River virus (L:AF234534; N:AAC54627); BBOV, Bimbo virus (L:GU816016); BEFV, bovine ephemeral fever virus (L:NP.065409; N:NP.065398); BGNV, Bangoran virus (L:GU816010); BRMV, Berrimah virus (L:AZ43265); BTKV, Boteke virus (L:GU816014); BYSMV, barley yellow striate mosaic virus (L:ACT21686); CFRV, China fish rhabdovirus (L:AAX86686); CHPV, Chandipura virus (L:P13179; N:P11211); CHVV9824, Charleville virus (L:AY854644); CHVV9847, Charleville virus (L:AY854672); COCV, Cocal virus (L:ACB47438; N:ACB47434); CPV, coastal plains virus (L:ADG86364; N:ADG86356); DAffSV, Drosophila affinis sigma virus 10 (L:GQ410980); DobsSV, Drosophila obscura sigma virus 10A (N:GQ410979); DURV, Durham virus (L:ADB88761; N:ADB88758); DUVV, Duvenhage virus (L:ABZ81216; N:Q66453); EBLV1, european bat lyssavirus 1 (L:ABZ81181; N:AAX62875); EBLV2, european bat lyssavirus 2 (L:ABZ81191; N:YP.001285393); FLAV, Flanders virus (L:AAN73288; N:AAN73283); FUKV, Fukuoka virus (L:AAZ43279); GARV, Garba virus (L:GU816018); HDOOV, Humpty doo virus (L:AAZ43271); HIRRV, Hirame rhabdovirus (L:NP_919035; N:NP_919030); IHNV, infectious hematopoietic necrosis virus (L:CAA52076; N:NP042676); IRKV, Irkut virus (L:ABV03823; N:Q5VKP6); ISFV, Isfahan virus (L:Q5K2K3; N:Q5K2K7); KAMV, Kamese virus (L:GU816011); KCV, Kern Canyon virus (N:ABE69215); KEUV, Keuraliba virus (L:GU816021); KHUV, Khujand virus (L:ABV03824; N:Q6X1D4); KIMV, Kimberley virus (L:AAZ43266); KOLV, Kolongo virus (L:GU816020; N:ABE69214); KOTV, Kotonkan virus (L:AAZ43267; N:ABE69213); LBV, Lagos bat virus (L:ABZ81171; N:ABF56214); LDV, Le Dantec virus (L:AAZ43278); LYMoV, lettuce yellow mottle virus (L:YP_002308376; N:YP_00208371); LNYV, lettuce necrotic yellows virus (L:YP-425092; N:CAG34083); MEBV, mount Elgon bat virus (N:ABE69217); MFSV, maize fine streak virus (L:YP-052849; N:YP-052843); MMV, maize mosaic virus (L:YP_052855; N:YP_052850); MOKV, Mokola virus (L:ABZ81211; N:YP_142350); MOSV, Mossuril virus (L:GU816012); MOUV, Moussa virus (L:ACZ81402; N:ACZ81403); NASV, Nasoule virus (L:GU816017); NCMV, northern cereal mosaic virus (L:NP_597914; N:NP_057954); NGAV, Ngaingan virus (L:YP_003518294; N:YP_003518280); NKOV, Nkolbisson virus (L:GU816022); OBOV, Obodhiang virus (N:ABE69212); OFV, orchid fleck virus (L:YP_001294929; N:BAH97109); OITAV, Oita rhabdovirus (N:BAD13431); OUAV, Ouango virus (L:GU816015); PCRV, Parry Creek virus (L:AAZ43275); PERV, Perinet virus (L:AAZ43280); PFRV, Pike fry rhabdovirus (L:ACP28002; N:ACP27998); PIRVV, Piry virus (N:P26037); PORV, Porton virus (L:GU816013); PYDV, potato yellow dwarf virus (N:ABW35154); RABV, Rabies virus (L:Q66T60; N:ACN15666); RBUV, Rochambeau virus (N:ABE69218); RYSV, rice yellow stunt virus (L:NP.620502; N:NP.620496); SCRV, Siniperca chuatsi rhabdovirus (L:YP.802942; N:YP.802937); SCV, strawberry crinkle virus (L:AAP03645); SFRV, Starry flounder rhabdovirus (L:AY450644); SHRV, snakehead rhabdovirus (L:NP_050585; N:NP_050580); SIGMAV, Drosophila melanogaster sigma virus (N:ACV67011); SIGMAVHAP23, Drosophila melanogaster sigma virus HAP23 (L:GQ375258; N:GQ375258); SIGMAVHAP30, Drosophila melanogaster sigma virus AP30 (L:YP_003126913; N:YP_003126908); SJAV, Sandjimba virus (L:GU816019; N:ABE69216); STRV, sea trout rhabdovirus (N:AAL35756); SVCV, spring viraemia of carp virus (L:Q91DR9; N:ABW24033); SYNV, Sonchus yellow net virus (L:NP_042286; N:NP_042281); TaVCV, Taro vein chlorosis virus (L:YP_224083; N:YP_224078); TIBV, Tibrogargan virus (L:ADG86355; N:ADG86347); TUPV, Tupaia virus (L:YP.238534; N:YP.238528); VHSV, viral hemorrhagic septicemia virus (L:CAB40833; N:P24378); VSAV, vesicular stomatitis Alagoas virus (L:ACB47443; N:ACB47439); VSIV, vesicular stomatitis Indiana virus (L:NP_041716; N:P11212); VSNJV, vesicular stomatitis New Jersey virus (L:P16379; N:P04881); WCBV, West Caucasian bat virus (L:ABV03821; N:Q5VKP2); WOGV, Wongabel virus (L:AAZ43276; N:YP_002333271).

1990; Muller and Standfast, 1986). In addition, OVRV was also isolated in 1984 from *Aedes vigilax* mosquitoes collected in Darwin (Weir et al., 1997). Taken together, OVRV strains have been isolated from disparate regions of tropical Australia, suggesting that this virus has been circulating widely for more than a decade.

404

405

406

407

408

An. annulipes s.l. is a species complex distributed across Australia, whilst *Ae. vigilax* has a coastal distribution and *Cx. edwardsi* is restricted to Queensland and the northern coast of New South Wales (Lee et al., 1984, 1987, 1989). Laboratory-based vector competence studies will be required to confirm whether

Please cite this article in press as: Quan, P.-L., et al., Genetic characterization of K13965, a strain of Oak Vale virus from Western Australia. Virus Res. (2011), doi:10.1016/j.virusres.2011.06.021

ARTICLE IN PRESS

P.-L. Quan et al. / Virus Research xxx (2011) xxx-xxx

mosquitoes of these species are vectors for OVRV. Blood meal stud-414 ies suggest that An. annulipes s.l. and Ae. vigilax feed on a variety 415 of vertebrate hosts, including birds, marsupials and cattle (Jansen 416 et al., 2009; Johansen et al., 2009b; Lee et al., 1984, 1987), whereas 417 Cx. edwardsi appear to prefer humans and birds (Lee et al., 1989). 418 No serosurveys of antibodies to K13965 in humans or animals in 419 the Ord River area have been conducted, and no antibodies to 420 CSIRO-1342 were found in cattle sera at Peachester (Cybinski and 421 Muller, 1990). OVRV likely cycles between mosquitoes and verte-422 brate host(s), possibly birds and/or pigs, as suggested by a reported 423 detection of antibodies to OVRV in feral pigs (Bourhy et al., 2008).

424 Phylogenetic analysis of the block III L aa sequence indicates 425 that OVRV is closely related to members of a recently proposed 426 Sandjimba group (Dacheux et al., 2010) within the tentative 427 Dimarhabdovirus supergroup (Bourhy et al., 2005) of the Rhab-428 doviridae family. Other members of this group were isolated from 429 birds or mosquitoes from the Central African Republic. However, 430 the Australian viruses K13965 and CSIRO-1342 form a branch 431 separate from the African viruses, likely reflecting independent 432 evolutionary pathways consistent with their geographic separa-433 tion. In this regard it is of interest that the Sandjimba group 434 435 forms a sister clade with the proposed Almpiwar group (Fig. 3), whose four members have been isolated from southern Queensland (Charleville virus strain Ch9824 from Charleville), from northern 437 Queensland (Almpiwar virus and Charleville virus strain Ch9847 438 from Mitchell River), and from the Northern Territory (Humpty Doo 130 virus from Beatrice Hill), similar geographic areas as Peachester, 440 Darwin and the Kimberley region where OVRV and K13965 have 441 been found (Fig. 1) (Bourhy et al., 2008). Thus, OVRV and Almpiwar 442 group viruses have shared or overlapping geographic distributions. 443 Analysis of the N aa sequence showed a similar relationship of OVRV 444 to two viruses from the Sandjima group for which N sequence is 445 available, and also indicated a relationship to TUPV and DURV that 446 was corroborated by the long L sequence analysis; however, fur-447 ther analyses are hampered by the lack of long L and N sequences 448 for other viruses, especially for Almpiwar group viruses. Additional 449 genomic sequence information will be necessary to provide further 450 insights into the evolutionary history and relationships of members 451 of these proposed groups. 452

Arbovirus surveillance in northern Australia has yielded insights 453 into the incidence, geographic range and ecology of major human 454 flavi- and alphaviral pathogens through continued annual studies 455 over the last 5 decades (Broom et al., 1998; Johansen et al., 2009a,b; 456 Liehne et al., 1976). Our identification of Oak Vale rhabdovirus 457 in uncharacterized isolates identified through those surveillance 458 efforts further defines viral flora in that area and enhances our 459 knowledge of the diversity and complexity of rhabdoviral genomes 460 and their phylogenetic relationships. 461

462 Acknowledgments

The authors are grateful to Michael D.A. Lindsay (Arbovirus 463 Surveillance and Research Laboratory, the University of West-464 ern Australia, Nedlands, Western Australia), Annette K. Broom 465 (Arbovirus Surveillance and Research Laboratory, the Univer-466 sity of Western Australia, Nedlands, Western Australia), and 467 Tony E. Wright (Mosquito Borne Disease Control Branch, 468 Western Australian Department of Health, Claremont, West-469 ern Australia) for mosquito collection, and technical staff in 470 the Arbovirus Surveillance and Research Laboratory for virus 471 isolation. We also thank Craig Street for bioinformatic analy-472 ses. This work was supported by National Institutes of Health 473 award AI57158 (Northeast Biodefense Center-Lipkin), contract 474 475 HHSN272201000040I/HHSN27200004/DO4, the United States 476 Agency for International Development's (USAID) Emerging Pandemic Threats (EPT) Program, PREDICT project, under terms of Cooperative Agreement Number GHN-A-OO-09-00010-00, the United States Department of Defense and The Western Australian Department of Health by funding the Arbovirus Surveillance and Research Program.

Appendix A. Supplementary data

Supplementary data associated with this article can be found, in the online version, at doi:10.1016/j.virusres.2011.06.021.

References

- Allison, A.B., Palacios, G., Travassos da Rosa, A., Popov, V.L., Lu, L., Xiao, S.Y., Detoy, K., Briese, T., Lipkin, W.I., Keel, M.K., Stallknecht, D.E., Bishop, G.R., Tesh, R.B., 2010. Characterization of Durham virus, a novel rhabdovirus that encodes both a C and SH protein. Virus Res. 155 (1), 112–122.
- Altschul, S.F., Gish, W., Miller, W., Myers, E.W., Lipman, D.J., 1990. Basic local alignment search tool. J. Mol. Biol. 215 (3), 403–410.
- Bellini, W.J., Englund, G., Rozenblatt, S., Arnheiter, H., Richardson, C.D., 1985. Measles virus P gene codes for two proteins. J. Virol. 53 (3), 908–919.
- Bourhy, H., Cowley, J.A., Larrous, F., Holmes, E.C., Walker, P.J., 2005. Phylogenetic relationships among rhabdoviruses inferred using the L polymerase gene. J. Gen. Virol. 86 (Pt 10), 2849–2858.
- Bourhy, H., Gubala, A.J., Weir, R.P., Boyle, D.B., 2008. Animal rhabdoviruses. In: Mahy, B.W.J.V.R., Van Regenmortel, M.H.V. (Eds.), Encyclopedia of Virology. , 3rd edn, pp. 111–121.
- Broom, A.K., Hall, R.A., Johansen, C.A., Oliveira, N., Howard, M.A., Lindsay, M.D., Kay, B.H., Mackenzie, J.S., 1998. Identification of Australian arboviruses in inoculated cell cultures using monoclonal antibodies in ELISA. Pathology 30 (3), 286–288.
- Broom, A.K., Lindsay, M., van Heuzen, B., Wright, T., Mackenzie, J., Smith, D., 1997. Contrasting patterns of flavivirus activity in the Kimberley region of Western Australia, 1992–1996. Arbovirus Res. Aust. (7), 25–30.
- Broom, A.K., Lindsay, M.D., Harrington, S.A., Smith, D.W., 2002. Investigation of the southern limits of Murray Valley encephalitis activity in Western Australia during the 2000 wet season. Vector Borne Zoonotic Dis. 2 (2), 87–95.
- Broom, A.K., Lindsay, M.D., Wright, A.E., Smith, D.W., Mackenzie, J.S., 2003. Epizootic activity of Murray Valley encephalitis and Kunjin viruses in an aboriginal community in the southeast Kimberley region of Western Australia: results of mosquito fauna and virus isolation studies. Am. J. Trop. Med. Hyg. 69 (3), 277–283.
- Calisher, C.H., Karabatsos, N., Zeller, H., Digoutte, J.P., Tesh, R.B., Shope, R.E., Travassos da Rosa, A.P., St George, T.D., 1989. Antigenic relationships among rhabdoviruses from vertebrates and hematophagous arthropods. Intervirology 30 (5), 241–257.
- Coll, J.M., 1995. The glycoprotein G of rhabdoviruses. Arch. Virol. 140 (5), 827–851.
- Crysler, J.G., Lee, P., Reinders, M., Prevec, L., 1990. The sequence of the nucleocapsid protein (N) gene of Piry virus: possible domains in the N protein of vesiculoviruses. J. Gen, Virol. 71 (Pt 9), 2191–2194.
- Curran, J., Kolakofsky, D., 2008. Nonsegmented negative-strand RNA virus RNA synthesis in vivo. Virology 371 (2), 227–230.
- Cybinski, D.H., Muller, M.J., 1990. Isolation of arboviruses from cattlee and insects at two sentinel sites in Queensland, Australia, 1979–85. Aust. J. Zool. 38, 25–32.
- Dacheux, L., Berthet, N., Dissard, G., Holmes, E.C., Delmas, O., Larrous, F., Guigon, G., Dickinson, P., Faye, O., Sall, A.A., Old, I.G., Kong, K., Kennedy, G.C., Manuguerra, J.C., Cole, S.T., Caro, V., Gessain, A., Bourhy, H., 2010. Application of broadspectrum resequencing microarray for genotyping rhabdoviruses. J. Virol. 84 (18), 9557–9574.
- Dietzgen, R.G., Callaghan, B., Wetzel, T., Dale, J.L., 2006. Completion of the genome sequence of Lettuce necrotic yellows virus, type species of the genus *Cytorhab- dovirus*. Virus Res. 118 (1–2), 16–22.
- Edgar, R.C., 2004. MUSCLE: multiple sequence alignment with high accuracy and high throughput. Nucleic Acids Res. 32 (5), 1792–1797.
- Elango, N., Kovamees, J., Varsanyi, T.M., Norrby, E., 1989. mRNA sequence and deduced amino acid sequence of the mumps virus small hydrophobic protein gene. J. Virol. 63 (3), 1413–1415.
- Gubala, A., Davis, S., Weir, R., Melville, L., Cowled, C., Walker, P., Boyle, D., 2010. Ngaingan virus, a macropod-associated rhabdovirus, contains a second glycoprotein gene and seven novel open reading frames. Virology 399 (1), 98–108.
- Gubala, A.J., Proll, D.F., Barnard, R.T., Cowled, C.J., Crameri, S.G., Hyatt, A.D., Boyle, D.B., 2008. Genomic characterisation of Wongabel virus reveals novel genes within the *Rhabdoviridae*. Virology 376 (1), 13–23.
- Huang, Y., Zhao, H., Luo, Z., Chen, X., Fang, R.X., 2003. Novel structure of the genome of Rice yellow stunt virus: identification of the gene 6-encoded virion protein. J. Gen. Virol. 84 (Pt 8), 2259–2264.
- Huelsenbeck, J.P., Ronquist, F., 2001. MRBAYES: Bayesian inference of phylogenetic trees. Bioinformatics 17 (8), 754–755.
- Jansen, C.C., Webb, C.E., Graham, G.C., Craig, S.B., Zborowski, P., Ritchie, S.A., Russell, R.C., van den Hurk, A.F., 2009. Blood sources of mosquitoes collected from urban and peri-urban environments in Eastern Australia with species-specific molecular analysis of avian blood meals. Am. J. Trop. Med. Hyg. 81 (5), 849–857.

Please cite this article in press as: Quan, P.-L., et al., Genetic characterization of K13965, a strain of Oak Vale virus from Western Australia. Virus Res. (2011), doi:10.1016/j.virusres.2011.06.021

7

482 483 484

477

478

479

480

481

485 486

487

488

489

490

491

492

493

494

495

496

497

498

499

500

501

502

503

504

505

506

507

508

509

510

511

512

513

514

515

516

517

518

519

520

521

522

523

524

525

526

527

528

529

530

531

532

533

534

535

536

537

538

539

540

541

542

543

544

545

546

547

548

549

550

55

552

G Model VIRUS 95485 1-8

8

553

554

555

556

557

558

559

560

561

562

563

564

565

566

567

568

569

570

571 572

573

574

575

576

577

578

579

580

581

582

583

584

585

586

587

588

589

590

591

592

593

594

595

596

597

598

599

600

601

602

603

604

605

606

607

608

609

610

611

612

613

614

615

616

617

618

619

620

621

622

623

624

625

626

627

628

P.-L. Quan et al. / Virus Research xxx (2011) xxx-xxx

- Jayakar, H.R., Whitt, M.A., 2002. Identification of two additional translation products from the matrix (M) gene that contribute to vesicular stomatitis virus cytopathology. J. Virol. 76 (16), 8011-8018.
- Johansen, C., Broom, A., Lindsay, M., Avery, V., Power, S., Dixon, G., Sturrock, K., Maley, F., McFall, S., Geerlings, K., Zammit, C., Masters, L., Bestall, A., Smith, D., 2009a. Arbovirus and vector surveillance in Western Australia, 2004/05 to 2007/08. Arbovirus Res. Aust. 10, 76-81.
- Johansen, C.A., Power, S.L., Broom, A.K., 2009b. Determination of mosquito (Diptera: Culicidae) bloodmeal sources in Western Australia: implications for arbovirus transmission. J. Med. Entomol. 46 (5), 1167-1175.
- Johansen, C.A., van den Hurk, A.F., Ritchie, S.A., Zborowski, P., Nisbet, D.J., Paru, R., Bockarie, M.J., Macdonald, J., Drew, A.C., Khromykh, T.I., Mackenzie, J.S., 2000. Isolation of Japanese encephalitis virus from mosquitoes (Diptera: Culicidae) collected in the Western Province of Papua New Guinea, 1997-1998. Am. J. Trop. Med. Hyg. 62 (5), 631-638.
- Kretzschmar, E., Peluso, R., Schnell, M.J., Whitt, M.A., Rose, J.K., 1996. Normal replication of vesicular stomatitis virus without C proteins. Virology 216 (2), 309-316.
- Kuzmin, I.V., Hughes, G.J., Rupprecht, C.E., 2006. Phylogenetic relationships of seven previously unclassified viruses within the family Rhabdoviridae using partial nucleoprotein gene sequences. J. Gen. Virol. 87 (Pt 8), 2323–2331.
- Kuzmin, I.V., Novella, I.S., Dietzgen, R.G., Padhi, A., Rupprecht, C.E., 2009. The rhabdoviruses: biodiversity, phylogenetics, and evolution. Infect. Genet. Evol. 9 (4), 541-553.
- Lee, D.J., Hicks, M.M., Debenham, M.L., Griffiths, M., Marks, E.N., Bryan, J.H., Russell, R.C., 1989. The Culicidae of the Australasian Region, Entomology Monograph No. 2, vol. 7. Australian Government Publishing Service, Canberra, Australia.
- Lee, D.J., Hicks, M.M., Griffiths, M., Debenham, M.L., Bryan, J.H., Russell, R.C., Geary, M., 1987. The Culicidae of the Australasian Region, Entomology Monograph No. 2, vol. 7. Australian Government Publishing Service, Canberra, Australia.
- Lee, D.J., Hicks, M.M., Griffiths, M., Russell, R.C., Marks, E.N., 1984. The Culicidae of the Australasian Region, Entomology Monograph No. 2, vol. 7. Australian Government Publishing Service, Canberra, Australia.
- Li, Z., Xu, J., Patel, J., Fuentes, S., Lin, Y., Anderson, D., Sakamoto, K., Wang, L.F., He, B., 2011. Function of the small hydrophobic protein of J paramyxovirus. J. Virol. 85 (1), 32-42.
- Liehne, P.F., Anderson, S., Stanley, N.F., Liehne, C.G., Wright, A.E., Chan, K.H., Leivers, S., Britten, D.K., Hamilton, N.P., 1981. Isolation of Murray Valley encephalitis virus and other arboviruses in the Ord River Valley 1972-1976. Aust. J. Exp. Biol. Med. Sci. 59 (Pt 3), 347-356.
- Liehne, P.F., Stanley, N.F., Alpers, M.P., Liehne, C.G., 1976. Ord River arboviruses-the study site and mosquitoes. Aust. J. Exp. Biol. Med. Sci. 54 (5), 487-497.
- Lindsay, M.D., Broom, A.K., Wright, A.E., Johansen, C.A., Mackenzie, J.S., 1993. Ross River virus isolations from mosquitoes in arid regions of Western Australia: implication of vertical transmission as a means of persistence of the virus. Am. I. Trop. Med. Hvg. 49 (6), 686-696.
- Longdon, B., Obbard, D.J., Jiggins, F.M., 2010. Sigma viruses from three species of Drosophila form a major new clade in the rhabdovirus phylogeny. Proc. Biol. Sci. 277 (1678), 35-44.
- Mackenzie LS Broom AK 1999 Ord river irrigation area: the effect of dam construction and irrigation on the incidence of Murray Valley encephalitis virus. In: Kay, B.H. (Ed.), Water Resources-Health, Environment and Development. Spon, London, pp. 108-122.
- Margulies, M., Egholm, M., Altman, W.E., Attiya, S., Bader, J.S., Bemben, L.A., Berka, J., Braverman, M.S., Chen, Y.J., Chen, Z., Dewell, S.B., Du, L., Fierro, J.M., Gomes, X.V., Godwin, B.C., He, W., Helgesen, S., Ho, C.H., Irzyk, G.P., Jando, S.C., Alenquer, M.L., Jarvie, T.P., Jirage, K.B., Kim, J.B., Knight, J.R., Lanza, J.R., Leamon, J.H., Lefkowitz, S.M., Lei, M., Li, J., Lohman, K.L., Lu, H., Makhijani, V.B., McDade, K.E., McKenna, M.P., Myers, E.W., Nickerson, E., Nobile, J.R., Plant, R., Puc, B.P., Ronan, M.T., Roth, G.T., Sarkis, G.J., Simons, J.F., Simpson, J.W., Srinivasan, M., Tartaro, K.R., Tomasz, A., Vogt, K.A., Volkmer, G.A., Wang, S.H., Wang, Y., Weiner, M.P., Yu, P., Begley, R.F., Rothberg, J.M., 2005. Genome sequencing in microfabricated high-density picolitre reactors. Nature 437 (7057), 376-380.
- McWilliam, S.M., Kongsuwan, K., Cowley, J.A., Byrne, K.A., Walker, P.J., 1997. Genome organization and transcription strategy in the complex GNS-L intergenic region of bovine ephemeral fever rhabdovirus. J. Gen. Virol. 78 (6), 1309-1317.
- Morzunov, S.P., Winton, J.R., Nichol, S.T., 1995. The complete genome structure and phylogenetic relationship of infectious hematopoietic necrosis virus. Virus Res. 38 (2-3), 175-192.
- Muller, M.J., Standfast, H.A., 1986. Vectors of ephemeral fever group viruses. In: St George, T.D., Kay, B.H., Blok, J. (Eds.), Arbovirus Research in Australia: Proceedings of the Fourth Symposium. CSIRO/QMIR, Brisbane, pp. 295-300.
- Nagai, Y., Kato, A., 2004. Accessory genes of the paramyxoviridae, a large family of nonsegmented negative-strand RNA viruses, as a focus of active investigation by reverse genetics. Curr. Top. Microbiol. Immunol. 283, 197-248.
- Nishizawa, T., Kurath, G., Winton, J.R., 1997. Sequence analysis and expression of the M1 and M2 matrix protein genes of hirame rhabdovirus (HIRRV). Dis. Aquat. Organ. 31, 9-17.

- Peluso, R.W., Richardson, J.C., Talon, J., Lock, M., 1996. Identification of a set of proteins (C' and C) encoded by the bicistronic P gene of the Indiana serotype of vesicular stomatitis virus and analysis of their effect on transcription by the viral RNA polymerase. Virology 218 (2), 335-342.
- Poch, O., Blumberg, B.M., Bougueleret, L., Tordo, N., 1990. Sequence comparison of five polymerases (L proteins) of unsegmented negative-strand RNA viruses: theoretical assignment of functional domains. J. Gen. Virol. 71 (Pt 5), 1153-1162.
- Poch, O., Sauvaget, I., Delarue, M., Tordo, N., 1989. Identification of four conserved motifs among the RNA-dependent polymerase encoding elements. Embo J. 8 (12), 3867-3874.
- Quan, P.L., Junglen, S., Tashmukhamedova, A., Conlan, S., Hutchison, S.K., Kurth, A., Ellerbrok, H., Egholm, M., Briese, T., Leendertz, F.H., Lipkin, W.I., 2010. Moussa virus: a new member of the Rhabdoviridae family isolated from Culex decens mosquitoes in Cote d'Ivoire. Virus Res. 147 (1), 17-24.
- Quan, P.L., Palacios, G., Jabado, O.J., Conlan, S., Hirschberg, D.L., Pozo, F., Jack, P.J., Cisterna, D., Renwick, N., Hui, J., Drysdale, A., Amos-Ritchie, R., Baumeister, E., Savy, V., Lager, K.M., Richt, J.A., Boyle, D.B., Garcia-Sastre, A., Casas, I., Perez-Brena, P., Briese, T., Lipkin, W.I., 2007. Detection of respiratory viruses and subtype identification of influenza A viruses by GreeneChipResp oligonucleotide microarray. J. Clin. Microbiol. 45 (8), 2359-2364.
- Reed, S.E., Tsai, C.W., Willie, K.J., Redinbaugh, M.G., Hogenhout, S.A., 2005. Shotgun sequencing of the negative-sense RNA genome of the rhabdovirus Maize mosaic virus. J. Virol. Methods 129 (1), 91-96.
- Revill, P., Trinh, X., Dale, J., Harding, R., 2005. Taro vein chlorosis virus: characterization and variability of a new nucleorhabdovirus. J. Gen. Virol. 86 (Pt 2), 491-499
- Rice, P., Longden, I., Bleasby, A., 2000. EMBOSS: the European Molecular Biology Open Software Suite. Trends Genet. 16 (6), 276-277.
- Scholthof, K.B., Hillman, B.I., Modrell, B., Heaton, L.A., Jackson, A.O., 1994. Characterization and detection of sc4: a sixth gene encoded by sonchus yellow net virus. Virology 204 (1), 279-288.
- Schutze, H., Mundt, E., Mettenleiter, T.C., 1999. Complete genomic sequence of viral hemorrhagic septicemia virus, a fish rhabdovirus. Virus Genes 19 (1), 59-65.
- Spiropoulou, C.F., Nichol, S.T., 1993. A small highly basic protein is encoded in overlapping frame within the P gene of vesicular stomatitis virus. J. Virol. 67 (6), 3103-3110.
- Springfeld, C., Darai, G., Cattaneo, R., 2005. Characterization of the Tupaia rhabdovirus genome reveals a long open reading frame overlapping with P and a novel gene encoding a small hydrophobic protein. J. Virol. 79 (11), 6781-6790.

Tamura, K., Dudley, J., Nei, M., Kumar, S., 2007. MEGA4: Molecular Evolution-

ary Genetics Analysis (MEGA) software version 4.0. Mol. Biol. Evol. 24 (8), 1596 - 1599

- Tanno, F., Nakatsu, A., Toriyama, S., Kojima, M., 2000. Complete nucleotide sequence of Northern cereal mosaic virus and its genome organization. Arch. Virol. 145 (7), 1373-1384.
- Tao, J.J., Zhou, G.Z., Gui, J.F., Zhang, Q.Y., 2008. Genomic sequence of mandarin fish rhabdovirus with an unusual small non-transcriptional ORF. Virus Res. 132 (1-2), 86-96.
- Teninges, D., Bras, F., Dezelee, S., 1993. Genome organization of the sigma rhab-
- dovirus: six genes and a gene overlap. Virology 193 (2), 1018–1023. Tordo, N., Benmansour, A., Calisher, C., Dietzgen, R.G., Fang, R.X., Jackson, A.O., Kurath, G., Nadin-Davis, S., Tesh, R.B., Walker, P.J., 2005. Virus Taxonomy. Eighth Report of the International Committee on Taxonomy of Viruses , pp. 623–644.
- Tordo, N., Poch, O., Ermine, A., Keith, G., Rougeon, F., 1986, Walking along the rabies genome: is the large G-L intergenic region a remnant gene? Proc. Natl. Acad. Sci. U.S.A. 83 (11), 3914-3918.
- Walker, P.J., Kongsuwan, K., 1999. Deduced structural model for animal rhabdovirus
- glycoproteins. J. Gen. Virol. 80 (Pt 5), 1211–1220. Wang, Y., McWilliam, S.M., Cowley, J.A., Walker, P.J., 1994. Complex genome organization in the GNS-L intergenic region of Adelaide River rhabdovirus. Virology 203 (1), 63-72.
- Weir, R.P., Hyatt, A.D., Calisher, C.H., Whelan, P.I., 1997. New records of arboviruses isolated from mosquitoes in the Northern Territory, 1982–1992. Arbovirus Res. Aust. 7, 311-321.
- Whelan, S., Goldman, N., 2001. A general empirical model of protein evolution derived from multiple protein families using a maximum-likelihood approach. Mol. Biol. Evol. 18 (5), 691-699.
- Whelan, S.P., Barr, J.N., Wertz, G.W., 2004. Transcription and replication of nonsegmented negative-strand RNA viruses. Curr. Top. Microbiol. Immunol. 283, 61 - 119
- Wilson, R.L., Fuentes, S.M., Wang, P., Taddeo, E.C., Klatt, A., Henderson, A.J., He, B., 2006. Function of small hydrophobic proteins of paramyxovirus. J. Virol. 80 (4), 1700-1709.
- Zhu, R.L., Lei, X.Y., Ke, F., Yuan, X.P., Zhang, Q.Y., 2011. Genome of turbot rhabdovirus exhibits unusual non-coding regions and an additional ORF that could be expressed in fish cell. Virus Res. 155 (2), 495-505.

629

630

Please cite this article in press as: Quan, P.-L., et al., Genetic characterization of K13965, a strain of Oak Vale virus from Western Australia. Virus Res. (2011), doi:10.1016/j.virusres.2011.06.021